arXiv:1301.1063v1 [math.DG] 6 Jan 2013

Castling transformations of projective structures

Hironao Kato *

December 11, 2013

Abstract

We investigate the castling transformations of prehomogeneous vector spaces from the view point of projective structures and Grassmannian structures. We give a certain classification of manifolds equipped with a flat projective structure obtained by a finite number of castling transformations.

1 Introduction

The classification of manifolds admitting a flat projective structure is still widely open (cf. [OT, chapter 6]) and active area. Indeed recently the paper [GC] was published and it is proved that a connected sum $\boldsymbol{R}P^3\#\boldsymbol{R}P^3$ does not admit a flat projective structure. In our last paper [Kat] we proved that invariant flat complex projective structures on complex Lie groups correspond to certain infinitesimal prehomogeneous vector spaces. In the theory of prehomogeneous vector spaces there is a notion of castling transformations, which is a certain transformation of linear representations of algebraic groups preserving the prehomogeneity. In this paper we try to establish a transformations of manifolds equipped with a projective structure as a generalization of castling transformations. As castling transformations preserve the prehomogeneity of representations, our castling transformations of projective structures preserve the projectively flatness. Moreover since we can repeat a castling transformation, we can construct a sequence of projectively flat manifolds from a given projectively flat manifold. In fact we prove the following: Let (Q, ω) be a flat Grassmannian Cartan connection of type (β, α) over a manifold M. Assume $\alpha + \beta \geq 3$ and $\alpha \leq \beta$.

Theorem 1.1. *By a finite number of castling transformations from (Q, ω) we obtain a projectively flat manifold N, which is a principal fiber bundle over M. There is a one-to-one correspondence between the set of the structure group $\prod PL(k_i)$ of N $(j \geq 1)$* and the set of solutions $(k_1, \ldots, k_j)$ with $k_i \geq \alpha$ for the Grassmannian type equation

$$(*)\quad \alpha\beta + k_1^2 + \cdots + k_j^2 - j - (\alpha+\beta)k_1 \cdots k_j + 1 = 0.$$

2010 *Mathematics subject classification.* Primary 53B10, 11S90; Secondary 53C10

Key words and phrases. projective structure; Grassmannian structure; prehomogeneous vector space.

[*] The author is supported by JSPS and JSPS Strategic Young Researcher Overseas Visits Program for Accelerating Brain Circulation.

Remark 1.2. Especially the case $\alpha = 1$ corresponds to the assumption that M admits a flat projective structure. Thus from a projectively flat manifold M we can obtain a projectively flat principal fiber bundle N over M with group $PL(k_1) \times \cdots \times PL(k_j)$ satisfying the equation $(*)$.

Remark 1.3. By Theorem 1.1 we can obtain a sequence of projectively flat manifolds, which are connected by manifolds equipped with a flat Grassmannian Cartan connection. Each projectively flat manifold N is right invariant under the group action of $PL(k_1) \times \cdots \times PL(k_j)$.

About the existence problem there is the following obstruction: A simply connected compact manifold admitting a flat projective structure is diffeomorphic to the sphere S^n (see [KN]). Thus the manifold $\prod_{i=1}^{j} S^{m_i}$ $(m_i, j \geq 2)$ does not admit any flat projective structure. This point distinguishes flat projective structures from flat affine connections and flat Riemannian metrics as any product of flat affine (resp. Riemannian) manifolds is flat affine (resp. Riemannian) manifold again. However in [Kat] we obtained a real Lie algebra $\mathfrak{sl}(k_1) \times \cdots \times \mathfrak{sl}(k_j)$ with a certain condition whose corresponding real Lie group admits a invariant flat real projective structure. The aim of this paper is also to generalize these examples from the view point of Grassmannian structures.

The paper is organized as follows. First of all we review the Grassmannian structures in § 1 and establish castling transformations of projective structures by using Cartan connections in § 2 and § 3. In § 4 we try to understand castling transformations from the view point of G-structures and linear connections. Indeed from a Grassmannian structure of type (n, m) on a manifold M we construct a linear connection $\widetilde{\chi}$ on a $PL(m)$-bundle M_m over M. As a result $\widetilde{\chi}$ is not torsion-free, however we see that there is a enough reason to call castling transformations of "projective structures". In § 5 with respect to the projective structure $[\widetilde{\chi}]$ on M_m, firstly we show that the fiber $PL(m)$ in M_m is totally geodesic and induces an invariant flat projective structure $[\nabla_m]$ on $PL(m)$. Since the base space M_m is locally trivial, we can naturally ask whether the product connection $\nabla \times \nabla'$ gives a projectively flat affine connection on the product manifolds. By using the Ricci tensor we see that $\nabla \times \nabla'_m$ is not projectively flat, where ∇ is any flat affine connection on M and ∇'_m is any affine connection on $PL(m)$ projectively equivalent to ∇_m. In § 6 we investigate the base spaces obtained by successive castling transformations and describe a relation between base spaces. § 7 is devoted to some examples of base spaces. In § 8 we investigate the positive integer solutions of the Grassmannian type equation $(*)$, and give one conjecture.

2 Preliminaries

2.1 Review of Grassmannian structures and projective structures

We recall the notion of Grassmannian structures and projective structures to establish castling transformations in the differential geometry. Let M be a real manifold of dimension r. Denote by $\mathcal{L}(M)$ a bundle of linear flames of M and we regard an element of $\mathcal{L}(M)$ as a linear isomorphism $\boldsymbol{R}^r \to T_pM$. We identify $\boldsymbol{R}^r$ with $\boldsymbol{R}^n \otimes R^m$ and consider a $GL(n) \otimes GL(m)$-structure P_tM, i.e. a subbundle of $\mathcal{L}(M)$ with structure group $GL(n) \otimes GL(m)$. If we have $n, m \geq 2$, we call P_tM a Grassmannian structure of type (n, m) in this paper. Note that if $n = 1$ or $m = 1$, then $P_tM = \mathcal{L}(M)$. Put $l := m + n$. There are various names and definition. In [Han] and [Ish] a $GL(n) \otimes GL(m)$-structure is called a tensor product structure. On the other hand in

[MS] an isomorphism $\sigma : TM \to V \otimes W$ itself is called a Grassmannian structure, where V and W are vector bundles with rank n and m over M $(n, m \geq 2)$. Such an isomorphism σ gives a $GL(n) \otimes GL(m)$-structure in a natural manner, however the author does not know whether the converse is true. Typical examples admitting a Grassmannian structure are Grassmannian manifolds (see [MS] for other examples). Denote by $Gr_{m,m+n}$ a Grassmannian manifold $Gr_{m,m+n}$ consisting of m-dimensional subspaces in the $(m+n)$-dimensional real vector space $\boldsymbol{R}^{m+n}$. The real projective transformation group $PL(\boldsymbol{R}^{m+n})$ acts on $Gr_{m,m+n}$ transitively. We fix a basis of $\boldsymbol{R}^{m+n}$. Then $PL(\boldsymbol{R}^{m+n})$ is expressed as the quotient $GL(m+n)/\boldsymbol{R}^* I_{m+n}$, which we denote by $PL(m+n)$. Let v be a linear frame $(e_1, \cdots, e_m)$ of W_0. We denote by $< v >$ the m-dimensional subspace spanned by v. Let $PL(m+n)_{<v>}$ be the isotropy subgroup at $<v>$. Then we have $Gr_{m,m+n} = PL(m+n)/PL(m+n)_{<v>}$. The Lie algebra of $PL(m+n)$ is isomorphic to $\mathfrak{sl}(m+n)$, which has the graded decomposition $\mathfrak{sl}(m+n) = \mathfrak{g}_{-1} + \mathfrak{g}_0 + \mathfrak{g}_1$ given by

$$\begin{aligned}
\mathfrak{g}_{-1} &= \left\{ \begin{pmatrix} 0 & 0 \\ C & 0 \end{pmatrix} \middle| \, C \in M(n,m) \right\}, \\
\mathfrak{g}_0 &= \left\{ \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \middle| \, \begin{array}{l} A \in \mathfrak{gl}(m), \ B \in \mathfrak{gl}(n) \\ \mathrm{tr}(A+B) = 0 \end{array} \right\}, \\
\mathfrak{g}_1 &= \left\{ \begin{pmatrix} 0 & D \\ 0 & 0 \end{pmatrix} \middle| \, D \in M(m,n) \right\}.
\end{aligned}$$

Let $\{e_\alpha\}$ be the natural basis of $\boldsymbol{R}^n$ and $\{f_i\}$ be the one of $\boldsymbol{R}^m$. We denote the dual basis of $\boldsymbol{R}^{n*}$ and $\boldsymbol{R}^{m*}$ by $\{e^\alpha\}$ and $\{f^i\}$ respectively. Then $\boldsymbol{R}^n \otimes \boldsymbol{R}^m$ is naturally identified with $\mathfrak{g}_{-1}$ and the isotropy representation $\rho : PL(m+n)_{<v>} \to GL(g_{-1})$ is given by

$$\rho : \begin{pmatrix} A & C \\ 0 & B \end{pmatrix} \mapsto B \otimes *A.$$

Thus the image of ρ is the group $GL(n) \otimes GL(m)$. Similarly $\boldsymbol{R}^{m*} \otimes \boldsymbol{R}^{n*}$ is identified with $\mathfrak{g}_1$. We write an element of $\boldsymbol{R}^{m*} \otimes \boldsymbol{R}^{n*}$ as $\sum_{\alpha_i} \eta_{\alpha_i} f^i \otimes e^\alpha$. We denote the vector $e_\alpha \otimes f_i$ by e_{α_i}. The isotropy representation ρ enables us to identify $GL(n) \otimes GL(m)$ with the subgroup G_0 of $PL(m+n)$;

$$G_0 = \left\{ \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \middle| \, A \in GL(m), B \in GL(n) \right\}.$$

Thus its Lie algebra $gl(n) \otimes I_m + I_n \otimes gl(m)$ is identified with $\mathfrak{g}_0$.

Let $P_t M$ be a $GL(n) \otimes GL(m)$-structure on M. The canonical form θ of $P_t M$ takes the value in $\boldsymbol{R}^n \otimes \boldsymbol{R}^m$. Following [Tan2] we explain equivalence classes of linear connections in $P_t M$. Let χ and χ' be connections in $P_t M$. Then $\chi' \sim \chi$ if and only if there is a $\boldsymbol{R}^{m*} \otimes \boldsymbol{R}^{n*}$-valued function F on $P_t M$ such that $\chi' - \chi = [\theta, F]$. Note that here we regard the value of θ, χ and F as an element of $\mathfrak{g}_{-1}$, $\mathfrak{g}_0$ and $\mathfrak{g}_1$. We call this equivalence relation a Grassmannian equivalence relation and denote the equivalence class of χ by $[\chi]$. Two equivalent connections have the same torsion. We consider the torsion tensor field as a map T: $P_t M \to \mathfrak{g}_1 \otimes \mathfrak{g}_1 \otimes \mathfrak{g}_{-1}$. Conventionally the linear map $x \circ T(x^{-1} \cdot, x^{-1} \cdot)$: $T_p M \times T_p M \to T_p M$ is called the torsion tensor. Moreover the tensor field T^*: $P_t M \to \mathfrak{g}_1 \otimes \mathfrak{g}_0$ is defined by

$$T^*(\xi) = \frac{1}{2l} \sum_{\alpha, i} [T_x(\xi, e_\alpha \otimes f_i), f^i \otimes e^\alpha] \qquad (\xi \in \mathfrak{g}_{-1}, x \in P_t M).$$

Then P_tM admits a unique equivalence class of connections $[\chi]$ such that $T^* = 0$, which is called an admissible class. From an admissible class $[\chi]$ on P_tM, we can construct a normal Cartan connection over M as follows. Let Q be the extended bundle of P_tM by the injection $\iota : G_o \hookrightarrow PL(m+n)_{<v>}$, i.e. $Q = P_tM \times_{G_o} PL(m+n)_{<v>}$. We denote the natural injection $P_tM \hookrightarrow Q$ by $\tilde{\iota}$. Put $l := m+n$. We define the $\mathfrak{sl}(m+n)$-valued 1 form ω on Q by

$$\begin{aligned} \omega(\tilde{\iota}_* X) &:= \theta(X) + \chi(X) + J_x(\theta(X)) \ \in \ \mathfrak{g}_{-1} + \mathfrak{g}_0 + \mathfrak{g}_1 \\ &= \begin{pmatrix} -{}^t\chi_m(X) & J_x(\theta(X)) \\ \theta(X) & \chi_n(X) \end{pmatrix} \quad (X \in T_xP_tM), \end{aligned} \tag{2.1}$$

where J is the tensor field $P_tM \ni x \mapsto J_x \in \mathfrak{g}_1 \otimes \mathfrak{g}_1$ of type $(0,2)$ on M defined by

$$J_x(e_{\alpha_i})_{\beta_j} = \frac{1}{(l^2-4)l}\{(2-l^2)\mathrm{Ric}_x(e_{\alpha_i}, e_{\beta_j}) - l\mathrm{Ric}_x(e_{\beta_i}, e_{\alpha_j}) - l\mathrm{Ric}_x(e_{\alpha_j}, e_{\beta_i}) - 2\mathrm{Ric}_x(e_{\beta_j}, e_{\alpha_i})\}. \tag{2.2}$$

We can obtain this expression of J by applying the technique of [Tan1, p.11] to [Tan2, Propsition 7.5]. Then the pair (Q, ω) gives a Grassmannian Cartan connection of type (n, m), thus by definition satisfies the following condition:

(1) ω: $T_uQ \to \mathfrak{sl}(m+n)$ gives a linear isomorphism,

(2) $R_g^*\omega = Ad(g^{-1})\omega$ for $g \in PL(m+n)_{<v>}$,

(3) $\omega(A^*) = A$ $(A \in \mathfrak{sl}(m+n)_{<v>})$, where A^* is the fundamental vector field.

The construction of Cartan connection ω depends on the choice of χ. However it is proved that if two linear connections in P_tM are equivalent, then the induced normal Grassmannian Cartan connections are isomorphic.

A $\mathfrak{g}$-valued 2-form Ω on Q defined by $\Omega = d\omega + \frac{1}{2}[\omega, \omega]$ is called a curvature form. A Grassmannian Cartan connection (Q, ω) is said to be flat if $\Omega = 0$. For an element $\xi \in \mathfrak{g}_{-1}$ we define a vector field $\check{\omega}(\xi)$ on Q by $\check{\omega}(\xi) := \omega^{-1}(\xi)$. A $\mathfrak{g}_0 + \mathfrak{g}_1$-valued 1-form Ω^* on Q defined by

$$\Omega_u^*(\xi) = \frac{1}{2l}\sum_{(\alpha,i)}[\Omega_u(\check{\omega}(\xi), \check{\omega}(e_\alpha \otimes f_i)), f^i \otimes e^\alpha] \quad (u \in Q, \xi \in \mathfrak{g}_{-1})$$

is called a $*$-curvature of (Q, ω) (see [Tan3]). A Grassmannian Cartan connection is called normal if $\Omega^* = 0$. In (2.1) the Grassmannian Cartan connection has been constructed from an admissible linear connection in P_tM, which is shown to be normal (see [Tan2]). Hence we obtain the following map

Φ: $\{ \ (P_tM, [\chi]) \mid [\chi]$ is an admissible class of linear connections in $P_tM \ \} \to \{$ normal Grassmannian Cartan connections on $M \ \}/\sim$.

The map Φ is shown to be bijective. We outline that Φ is surjective. Let (Q, ω) be a Grassmannian Cartan connection of type (n, m) on M. Let $\rho : PL(m+n)_{<v>} \to GL(g_{-1})$ be the isotropy representation. We denote the kernel of ρ by $ker\rho$. Then $PL(m+n)_{<v>}/ker\rho \cong GL(n) \otimes GL(m)$. Thus the quotient manifold $\widetilde{Q} := Q/ker\rho$ is regarded as a principal fiber bundle over M with structure group $GL(n) \otimes GL(m)$. Let ω_{-1} be the $\mathfrak{g}_{-1}$ component of 1-form ω. By using the natural projection $p : Q \to \widetilde{Q}$, we obtain the $\mathfrak{g}_{-1}$-valued 1-form θ on $\widetilde{Q}$ defined by $\rho^*\theta = \omega_{-1}$. Then $(\widetilde{Q},\ \theta)$ can be regarded as $GL(n) \otimes GL(m)$-structure and its canonical form. We define an injection $\iota : GL(n) \otimes GL(m) \hookrightarrow PL(l)_{<v>}$ by

$$\iota(A \otimes B) = \begin{pmatrix} {}^tB^{-1} & 0 \\ 0 & A \end{pmatrix}.$$

The homogeneous space $PL(l)_{<v>}/\iota(GL(n)\otimes GL(m))$ is homeomorphic to $\mathfrak{g}_1$, and hence there exists a bundle homomorphism $h:\widetilde{Q}\hookrightarrow Q$ corresponding to ι such that $\rho\circ h = id$. Then we can obtain the connection form χ defined by

$$\chi(X) = \omega(h_*X)_0 \quad (X\in TP_tM).$$

If (Q,ω) is normal, the induced connection χ satisfies $T^*=0$. Moreover $\Phi(\widetilde{Q},[\chi])$ coincides with the isomorphism class of (Q,ω). In this paper we call a pair $(P_tM,[\chi])$ a grassmannian structure (gr-structure) of type (n,m), where $[\chi]$ is an admissible class. This name may agree with the one in [Ish]. Especially when $m,n\neq 1$, $GL(n)\otimes GL(m)$-structure P_tM admits a unique admissible class of linear connections by Proposition 9.5 and Theorem 10.2 of [Tan2], and hence we obtain the following one-to-one correspondence (cf. [Tan2, Theorem 9.4]):

$\{GL(n)\otimes GL(m)$-structure of $M\}\to\{$normal Grassmannian Cartan connections over $M\}/_\sim$.

Now we consider the case $m=1$. Then the above discussion for the case $m=1$ yields the projective space $P(\boldsymbol{R}^{1+n})$, a grading of $\mathfrak{sl}(1+n)$, projective equivalence relation, a projective Cartan connection instead of Grassmannian manifolds $Gr_{m,m+n}$, the grading of $\mathfrak{sl}(m+n)$, Grassmannian equivalence relation, Grassmannian Cartan connections. For more details of projective Cartan connections we refer the reader to [Tan2], [Tan1] and [Aga]. From the appendix of [NS] we see that two linear connections are projectively equivalent iff they have the same torsion and the same set of pregeodesics. Conventionally a projective equivalence class of torsion-free linear connections in $\mathcal{L}(M)$ is called a projective structure. In the projective case the tensor T^* vanishes iff the torsion T vanishes. Thus an admissible class of linear connections on $\mathcal{L}(M)$ is a projective equivalence class consisting of torsion-free linear connections. It follows that a gr-structure $(P_tM,[\chi])$ for the case $m=1$ or $n=1$ is equivalent to a projective structure. Thus we have the following one-to-one correspondence:

$\{[\chi]:$ projective structures on $M\}\to\{$normal projective Cartan connections over $M\}/_\sim$.

A projective structure $[\chi]$ is said to be flat if χ is locally projectively equivalent to the standard flat affine connection χ_0. It is well known that χ is projectively flat iff χ is torsion-free and the projective curvature tensor of Weyl vanishes. Moreover a flat projective structure $[\chi]$ on M corresponds to a $(PL(1+n),P(\boldsymbol{R}^{1+n}))$-structure on M (see [Gol], [Kat] for the definition). Likewise a flat Grassmannian Cartan connection corresponds to a $(PL(m+n),Gr_{m,m+n})$-structure. The flatness of Grassmannian Cartan connection of type (n, m) $(n,m>2)$ should be described by the terminology of $GL(n)\otimes GL(m)$-structure. However the author does not know it. Let (Q,ω) be a projective Cartan connection over n-dimensional manifold. We also call (Q,ω) a Grassmannian Cartan connection of type $(n,1)$. Thus a normal Grassmannian Cartan connection of type (n,m) corresponds to a Grassmannian structure of type (n,m) if $n,m\geq 2$, and corresponds to a projective structure if $n=1$ or $m=1$.

2.2 Subgeometry

For the later argument, we introduce the notion of subgeometry, following [Gol]. Let A/B and A'/B' be real homogeneous spaces. We say that A/B is a subgeometry of A'/B' if there exists a Lie group homomorphism $F:A\to A'$ satisfying the following conditions:

(1) $F(B)\subset B'$,

(2) the induced map $\hat{F} : A/B \to A'/B'$ is a local diffeomorphism.

Let us denote the Lie algebra of A by $\mathfrak{a}$, the one of B by $\mathfrak{b}$. Likewise we define $\mathfrak{a}'$ and $\mathfrak{b}'$ for A' and B' respectively. Let Λ and Λ' be Maurer-Cartan forms of A and A'. Then F gives a bundle homomorphism corresponding to $F|_B : B \to B'$ and satisfies $F^*\Lambda' = dF \circ \Lambda$.

Proposition 2.1. *Let (Q, ω) be a Cartan connection of type A/B on M. Then there exists a Cartan connection (Q', ω') of type A'/B' on M.*

Proof. Since B acts on B' via F, from the given principal bundle Q we obtain the extended bundle $Q' = Q \times_B B'$. The bundle homomorphism $\widetilde{F} : Q \to Q'$ is defined by $u \mapsto [u, e]$, which corresponds to the restriction of F to B. Next we define a $\mathfrak{a}'$-valued 1-form ω' on $\widetilde{F}(Q)$ by

$$\omega'_{[u,e]}(\widetilde{F}_* X + Z^*) = dF \circ \omega(X) + Z \quad (X \in T_u Q, Z \in \mathfrak{b}').$$

We enlarge this definition to the whole of Q' by

$$\omega'_{[u,c]} = R^*_{c^{-1}} Ad(c^{-1}) \omega_{[u,e]} \quad (c \in B').$$

To verify that this definition is well defined, it is enough to check the following two cases:

(1) $\widetilde{F}_* X = Z^*_{[u,e]}$ $(X \in T_u Q,\ Z \in \mathfrak{b}')$. In this case there exists $Y \in \mathfrak{b}$ such that $X = Y^*_u$. Then $\widetilde{F}_* X = dF(Y)^*_{[u,e]}$. Hence

$$\begin{aligned} \omega'(\widetilde{F}_* X) &= dF \circ \omega(X) = dF(Y), \\ \omega'(Z^*_{[u,e]}) &= \omega'(dF(Y)^*_{[u,e]}) = dF(Y). \end{aligned}$$

Therefore $\omega'(\widetilde{F}_* X) = \omega'(Z^*_{[u,e]})$.

(2) $[ub, e] = [u, F(b)]$ $(u \in Q,\ b \in B)$. In this case we can verify $\omega'_{[ub,e]} = \omega'_{[u,F(b)]}$ by the following computation.

$$\begin{aligned} \omega_{[ub,e]}(\widetilde{F}_* X + Z^*_{[ub,e]}) &= dF \circ \omega_{ub}(X) + Z \\ &= dF(Ad(b^{-1}) \omega_u({R_{b^{-1}}}_* X)) + Z \\ &= dF(\omega_u(X)) + Z. \end{aligned}$$

On the other hand

$$\begin{aligned} \omega_{[u,e]}({R_{F(b)^{-1}}}_* \widetilde{F}_* X) &= \omega'_{[u,e]}(\widetilde{F}_* {R_{b^{-1}}}_* X) \\ &= dF \omega_u({R_{b^{-1}}}_* X) \\ &= dF(Ad(b) \omega_u(X)) \\ &= Ad(F(b)) dF(\omega_u(X)). \end{aligned}$$

$$\begin{aligned} Z^*_{[ub,e]} &= Z^*_{[u,F(b)]} \\ &= {R_{F(b)}}_*(Ad(F(b))Z)^*_{[u,e]}. \end{aligned}$$

Moreover we can verify (Q', ω') gives a Cartan connection of type A'/B' on M. □

Definition 2.2. *Let (Q, ω) and (Q', ω') be Cartan connections of type A/B and A'/B' respectively on M. Then we call (Q, ω) a subgeometry of (Q', ω') if there exists a bundle homomorphism $\iota : Q \to Q'$ corresponding to $F|_B : B \to B'$ such that ι induces the identity map between the base spaces and $\iota^* \omega' = dF \circ \omega$.*

In Proposition 2.1 a given Cartan connection (Q, ω) of type A/B induces (Q', ω') of type A'/B', and (Q, ω) is a subgeometry of (Q', ω').

Proposition 2.3. *Assume that a Cartan connection (Q, ω) is a subgeometry of (Q', ω'). If (Q, ω) is flat, then (Q', ω') is also flat. Moreover when the differential $dF : \mathfrak{a} \to \mathfrak{b}$ is an injective homomorphism, the converse is also true.*

Proof. We compute the curvature form Ω' of (Q', ω'). Pulling back Ω' by ι yields

$$\begin{aligned} \iota^*\Omega' &= \iota^*(d\omega' + \frac{1}{2}[\omega', \omega']) \\ &= dF(d\omega + \frac{1}{2}[\omega, \omega]) \\ &= dF(\Omega). \end{aligned}$$

Hence the assertion of the proposition follows. □

We fix the complementary subspace $\mathfrak{m}$ of $\mathfrak{b}$ and $\mathfrak{m}'$ of $\mathfrak{b}'$. Let ρ be the linear isotropy representation of B on the tangent space to A/B at the origin o. By identifying T_oA/B with $\mathfrak{m}$, ρ is given by $\rho(b)X = Ad(b)X + \mathfrak{b}$ for $b \in B$ and $X \in \mathfrak{m}$. Thus we obtain the two linear isotropy representations $\rho : B \to GL(\mathfrak{m})$ and $\rho' : B' \to GL(\mathfrak{m}')$. We denote the kernel of ρ by C and the one of ρ' by C'. Since we assume that A/B is a subgeometry of A'/B', there is a homomorphism $F : A \to A'$ whose differential dF induces the linear isomorphism $\widehat{dF} : \mathfrak{m} \to \mathfrak{m}'$.

Lemma 2.4. *There exists an injective homomorphism $\overline{F} : \rho(B) \to \rho'(B')$ defined by $\overline{F} : \rho(b) \mapsto \widehat{dF} \circ \rho(b) \circ \widehat{dF}^{-1}$, and we have the commutative diagram:*

$$\begin{tikzcd} B \arrow[r, "F"'] \arrow[d, "\rho"'] \arrow[dr, phantom, "\circlearrowleft"] & B' \arrow[d, "\rho'"'] \\ \rho(B) \arrow[r, hook, "\overline{F}"'] & \rho'(B') \end{tikzcd}$$

Moreover F is regarded as a bundle homomorphism corresponding to $F : C \to C'$.

Proof. First we verify $F(C) \subset C'$. Assume that $b \in C$. Then $\rho(b)(X + \mathfrak{b}) = X + \mathfrak{b}$ for $X \in \mathfrak{m}$. Then $\rho'(F(b))(\widehat{dF}(X)) = Ad(F(b))(dF(X) + \mathfrak{b}') = dF(Ad(b)X) + \mathfrak{b}' = dF(X) + \mathfrak{b}' = \widehat{dF}(X)$. Thus $\rho'(F(b)) = id_{\mathfrak{m}'}$, and $F(b) \in C'$. We define $\overline{F} : \rho(B) \to \rho'(B')$ by $\bar{F} : \rho(b) \mapsto \rho'(F(b))$ for $b \in B$. Since $F(C) \subset C'$, this is well defined, moreover we have $\overline{F}(\rho(b)) = \widehat{dF} \circ \rho(b) \circ \widehat{dF}^{-1}$. It follows that $\overline{F}$ is injective. □

Let (Q, ω) and (Q', ω') be Cartan connections of type A/B and A'/B' respectively. Assume that (Q, ω) is a subgeometry of (Q', ω'). We denote the quotient manifold Q/C by $\widetilde{Q}$ and Q'/C' by $\widetilde{Q}'$. Then we obtain $\rho(B)$-structure $(\widetilde{Q}, \theta)$ and $\rho'(B')$-structure $(\widetilde{Q}', \theta')$ (see [Tan3, p.136]). The projection $\rho : Q \to \widetilde{Q}$ is corresponding to $\rho : B \to \rho(B)$. Recall that $(\widetilde{Q}, \theta)$ gives a $\rho(B)$-structure on M as follows: concerning each point $\rho(u) \in \widetilde{Q}$, $\rho(u)^{-1}$ is regarded as a linear isomorphism $T_{\pi_{Q/C}(\rho(u))}M \to \mathfrak{m}$ by $\rho(u)^{-1} : {\pi_{Q/C}}_* \rho_* X \mapsto \theta(\rho_* X) = \omega_{\mathfrak{m}}(X)$. Hence we obtain the bundle homomorphism $\widetilde{Q} \hookrightarrow L(M)$ corresponding to the inclusion $\rho(B) \to GL(\mathfrak{m})$.

Likewise we obtain the map $\widetilde{Q}' \hookrightarrow L(M)$, where $L(M)$ is regarded as the set of all linear isomorphism $y : \mathfrak{m}' \to T_pM (p \in M)$.

The bundle homomorphism $t : L(M) \to L(M)$ is defined by $t : x \mapsto x \circ \hat{dF}^{-1}$, which is corresponding to $GL(\mathfrak{m}) \ni A \mapsto \hat{dF} \circ A \circ \hat{dF}^{-1} \in GL(\mathfrak{m}')$.

Proposition 2.5. *The $\rho(B)$-structure $\widetilde{Q}$ is a reduction of $\rho'(B')$-structure $\widetilde{Q}'$ i.e.* 1) *there exists a bundle homomorphism $\bar{\iota} : \widetilde{Q} \hookrightarrow \widetilde{Q'}$ corresponding to $\bar{F} : \rho(B) \to \rho'(B')$, and* 2) $\bar{\iota}^*\theta' = \hat{dF} \circ \theta$. *The injection $\bar{\iota}$ is given by the restriction of $t : L(M) \to L(M)$.*

Proof. From assumption (Q, ω) is a subgeometry of (Q', ω'), thus we have a bundle homomorphism $\iota : Q \to Q'$ corresponding to $F|_B : B \to B'$. We define a map $\bar{\iota} : \widetilde{Q} \to \widetilde{Q}'$ by $\bar{\iota} : \rho(u) \to \rho' \circ \iota(u)$. Since $F|_B$ is a bundle homomorphism corresponding to $F : C \to C'$, $\bar{\iota}$ is well-defined and gives a bundle homomorphism corresponding to $\bar{F} : \rho(B) \to \rho'(B')$. Hence we obtain the commutative diagram:

$$\begin{tikzcd}
Q \arrow[r, "i"'] \arrow[d, "\rho"'] \arrow[dr, phantom, "\circlearrowleft"] & Q' \arrow[d, "\rho'"] \\
\widetilde{Q} \arrow[r, hook, "\bar{\iota}"'] \arrow[d, "\pi_{\widetilde{Q}}"'] \arrow[dr, phantom, "\circlearrowleft"] & \widetilde{Q}' \arrow[d, "\pi_{\widetilde{Q}'}"] \\
M \arrow[r, "id"'] & M.
\end{tikzcd} \tag{2.3}$$

Since $\bar{\iota}$ induces the identity of base spaces and $\bar{F}$ is injective, $\bar{\iota}$ is injective. Now we show that $\bar{\iota}^*\theta' = \hat{dF} \circ \theta$. Since $\bar{\iota} \circ \rho = \rho' \circ \iota$, pulling back $\bar{\iota}^*\theta'$ by $\rho : Q \to \widetilde{Q}$ yields $\rho^*(\bar{\iota}^*\theta') = (\rho' \circ \iota)^*\theta' = \iota * \omega'_{\mathfrak{m}'} = \hat{dF} \circ \omega_{\mathfrak{m}}$. Hence $\bar{\iota}^*\theta' = \hat{dF} \circ \theta$.

By using the inclusion $\widetilde{Q} \hookrightarrow L(M)$ and $\widetilde{Q}' \hookrightarrow L(M)$, t and $\bar{\iota}$, we obtain the following diagram, which will be shown commutative as follows.

$$\begin{tikzcd}
L(M) \arrow[r, "t"'] \arrow[dr, phantom, "\circlearrowleft"] & L(M) \\
\widetilde{Q} \arrow[u] \arrow[r, hook, "\bar{\iota}"'] & \widetilde{Q}' \arrow[u]
\end{tikzcd} \tag{2.4}$$

From the equality $\bar{\iota}^*\theta' = \hat{dF} \circ \theta$, for $\rho(u) \in \widetilde{Q}$ we have $\theta'_{\bar{\iota}(\rho(u))}(\bar{\iota}_*\rho_*X) = \hat{dF} \circ (\rho_*X) = \hat{dF} \circ \omega_{\mathfrak{m}}(X)$. Thus $\bar{\iota}(\rho(u))$ gives a linear isomorphism $T_{\pi_{\widetilde{Q}'}(\bar{\iota}(\rho(u)))}M \to \mathfrak{m}'$ by $\bar{\iota}(\rho(u)) : \pi_{\widetilde{Q}'}{}_*\bar{\iota}_*\rho_*X \mapsto \hat{dF} \circ \omega_{\mathfrak{m}}(X)$. Since the diagram (2.3) is commutative, we have $\pi_{\widetilde{Q}'}{}_*\bar{\iota}_*\rho_*X = \pi_{\widetilde{Q}}{}_*\rho_*X$. Therefore $\bar{\iota}(\rho(u)) = \rho(u) \circ \hat{dF}^{-1}$. On the other hand $t \circ \rho(u) = \rho(u) \circ \hat{dF}^{-1}$, and hence the diagram (2.4) is commutative. $\square$

Let X' be a homogeneous space of A'. We choose B' as the isotropy subgroup at a point v in X. If we are given a subgroup $A \subset A'$, we can consider the isotropy subgroup A_v of A at v. Then A/A_v gives a subgeometry of A'/B'. Henceforth we say that A/B is a subgeometry of A'/B' if this condition is satisfied: A is a subgroup of A' and B is the isotropy subgroup at v.

3 Castling transformations

In this section we establish the castling transformations of projective structures. Let G be a subgroup of $PL(l)$. We consider the homomorphism $F : G \times PL(m) \hookrightarrow PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$ defined by $(g, A) \mapsto g \otimes A$. By F we regard $G \times PL(m)$ as a subgroup of $PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$. When we identify $\boldsymbol{R}^l \otimes \boldsymbol{R}^m$ with $\underbrace{\boldsymbol{R}^l \oplus \cdots \oplus \boldsymbol{R}^l}_{m}$, $G \times PL(m)$ acts on $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$ by $(g, A).v := gv^tA$ for $v = (v_1, \ldots, v_m) \in P(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$. Assume that $n = l - m \geq 0$. Denote by $V_{m,l}$ a projective Stiefel manifold, which consists of projective frames of m-dimensional subspaces of $\boldsymbol{R}^l$.

Proposition 3.1. *Let G be a subgroup of $PL(l)$, and v be a point in $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$. The rank of $v = (v_1, \ldots, v_m)$ is m and the homogeneous space $G/G_{<v>}$ is a subgeometry of $Gr_{m,l} = PL(l)/PL(l)_{<v>}$ if and only if $G \times PL(m)/G \times PL(m)_v$ is a subgeometry of the projective space $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^m) = PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)/PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)_v$ defined by $F : G \times PL(m) \hookrightarrow PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$.*

Proof. We can prove this proposition by showing that the following four assertions are equivalent.

(1) $G/G_{<v>}$ is a subgeometry of the Grassmannian manifold $PL(l)/PL(l)_{<v>}$.

(2) $G. <v>$ gives an open orbit in $G_{m,l}$.

(3) $G \times PL(m).v$ gives an open orbit in $V_{m,l}$.

(4) $G \times PL(m)/G \times PL(m)_v$ is a subgeometry of the projective space $PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)/PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)_v$.

The proof of (1) $\Leftrightarrow$ (2) is easy. To prove (2) $\Leftrightarrow$ (3), we consider the fiber bundle

$$\begin{tikzcd}
V_{m,l} \arrow[d, "\pi"] & PL(m) \arrow[l] \\
G_{m,l} &
\end{tikzcd}$$

The natural projection π is continuous and open map. It follows (2) $\Leftrightarrow$ (3). To prove (3) $\Leftrightarrow$ (4), we observe that the manifold $V_{m,l}$ is naturally imbedded into $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$ with respect to the relative topology, indeed $V_{m,l}$ is an open submanifold in $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$. Next we consider the assertion $(3)'$: $G \times PL(m).v$ gives an open orbit in $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$. If $(3)'$ holds true, then v must belong to $V_{m,l}$. Hence we have (3) $\Leftrightarrow$ $(3)'$. The proof of the equivalence $(3)' \Leftrightarrow (4)$ is same as the one of (1) $\Leftrightarrow$ (2). □

Now assume that $n = l - m \geq 1$. We fix a point v of $V_{m,l}$ and one $v^\perp$ of $V_{n,l}$ such that the subspace $< v^\perp >$ spanned by $v^\perp$ is orthogonal to the one $< v >$ spanned by v. Let us define the isomorphism $* : PL(l) \to PL(l)$ by $g \mapsto {}^tg^{-1}$. Then $*$ gives the isomorphism between $PL(l)_{<v>}$ and $PL(l)_{<v^\perp>}$. We denote the differential of $*$ by the same symbol, and we have $*(A) = -{}^tA$ for $A \in \mathfrak{sl}(l)$. Then $G. <v>$ gives an open orbit in $Gr_{m,l}$ if and only if $*G. <v^\perp>$ gives an open orbit of $Gr_{n,l}$. Hence by proposition 3.1 we obtain the following:

Proposition 3.2. (cf. [SK]) The group $G \times PL(m)$ admits an open orbit in $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$ if and only if $*G \times PL(l - m)$ admits an open orbit in $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^{l-m})$.

Strictly in [SK] the above proposition is proved in the algebraic group category. Let $F : G \to GL(\boldsymbol{C}^l)$ be a rational representation of G. Then originally the transformation

$$F \otimes id : G \times GL(m) \to GL(\boldsymbol{C}^l \otimes \boldsymbol{C}^m) \Leftrightarrow F^* \otimes id : G \times GL(l-m) \to GL(\boldsymbol{C}^l \otimes \boldsymbol{C}^{l-m})$$

is called a castling transformation. Now we define the castling transformation of Cartan connections. Denote by $\mathfrak{g}$ the Lie algebra of G. Let G be a subgroup of $PL(l)$.

Proposition 3.3. *Denote by Q a principal fiber bundle over M with structure group $G_{<v>}$ and by ω a $\mathfrak{g}$-valued 1-form on Q. Then the following are equivalent.*

1. *(Q, ω) gives a Cartan connection on M of type $G/G_{<v>}$, which is a subgeometry of $Gr_{m,l} = PL(l)/PL(l)_{<v>}$*
2. *$(Q, *\omega)$ gives a Cartan connection on M of type $*G/*G_{<v^\perp>}$, which is a subgeometry of $Gr_{n,l} = PL(l)/PL(l)_{<v^\perp>}$*
3. *$(Q \times PL(m), \omega \times \Lambda_1)$ gives a Cartan connection of type $G \times PL(m)/G \times PL(m)_v$, which is a subgeometry of the projective space $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$. This subgeometry is defined by the homomorphism $\imath : G \times PL(m) \ni (g, A) \mapsto g \otimes A \in PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$.*
4. *$(Q \times PL(n), *\omega \times \Lambda_1)$ gives a Cartan connection of type $*G \times PL(n)/*G \times PL(n)_{v^\perp}$, which is a subgeometry of the projective space $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^n)$.*

Moreover if one of the above Cartan connections is flat, then the other Cartan connections are also flat. When one of the conditions (1)-(4) is satisfied, the Cartan connection (Q, ω) induces a $GL(n) \otimes GL(m)$-structure P_tM on M. The base space M_m of $(Q \times PL(m), \omega \times \Lambda_1)$ is the quotient of P_tM by $GL(n) \otimes GL(1)$.

Remark 3.4. If the condition (1) is satisfied, the Cartan connection $(Q, *\omega)$ induces a $GL(m) \otimes GL(n)$-structure P_tM' and the base space M_n of $(Q \times PL(n), *\omega \times \Lambda_1)$ is the quotient of P_tM' by $GL(m) \otimes GL(1)$. In fact the base space M_n is naturally identified with $P_tM/GL(1) \otimes GL(m)$ by Proposition 4.1, which will be proved later.

Proof. (1) $\Leftrightarrow$ (2): We define the action of $*G_{<v^\perp>}$ on Q by $u \cdot g := u * g$ ($u \in Q$, $g \in *G_{<v^\perp>}$). Then the bundle Q is regarded also as a principal fiber bundle over M with structure group $*G_{<v^\perp>}$. Moreover we define a one-form $*\omega$ by the composite of $* : \mathfrak{sl}(l) \to \mathfrak{sl}(l)$ and ω. We can easily verify that the assertions 1 and 2 in the proposition are equivalent. On the other hand since $G/G_{<v>}$ is a subgeometry of $*G/*G_{<v^\perp>}$, the Cartan connection (Q, ω) induces a Cartan connection (Q_c, ω_c) of type $*G/*G_{<v^\perp>}$ by Proposition 2.1. This induced Cartan connection is isomorphic with $(Q, *\omega)$. Hence by Proposition 2.3 (Q, ω) is flat if and only if $(Q, *\omega)$ is flat.

(1) $\Leftrightarrow$ (3): Assume the assertion 1. Then $Q \times PL(m)$ is regarded as a principal fiber bundle over M with structure group $G_{<v>} \times PL(m)$. Since $G \times PL(m)_v$ is a closed subgroup of $G_{<v>} \times PL(m)$, we have the quotient $Q \times PL(m)/G \times PL(m)_v$ over which $Q \times PL(m)$ is regarded as a principal fiber bundle with structure group $G \times PL(m)_v$. Then we can directly check $(Q \times PL(m), \omega \times \Lambda_1)$ gives a Cartan connection of type $G \times PL(m)/G \times PL(m)_v$ on $Q \times PL(m)/G \times PL(m)_v$. By Proposition 3.1 $G \times PL(m)/G \times PL(m)_v$ is a subgeometry of the projective space $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$. Conversely we assume the assertion (3). Then $G/G_{<v>}$ is a subgeometry of $Gr(m, l) = PL(l)/PL(l)_{<v>}$. We can directly check (Q, ω) gives a Cartan connection of type $G/G_{<v>}$ on M .

(2) $\Leftrightarrow$ (4): The proof is same with one of (1) $\Leftrightarrow$ (3).

Next we prove the equivalence of flatness of the above Cartan connections. We have already proved this equivalence about the case (1) $\Leftrightarrow$ (2). We consider the case (1) $\Leftrightarrow$ (3). We first observe that $(d\omega + \frac{1}{2}[\omega, \omega], d\Lambda_1 + \frac{1}{2}[\Lambda_1, \Lambda_1])$ gives $\mathfrak{g} \times \mathfrak{sl}(m)$-valued 2-form on $Q \times PL(m)$. We can directly verify

$$d(\omega \times \Lambda_1) + \frac{1}{2}[\omega \times \Lambda_1, \omega \times \Lambda_1] = (d\omega + \frac{1}{2}[\omega, \omega], d\Lambda_1 + \frac{1}{2}[\Lambda_1, \Lambda_1]).$$

Note that since Λ_1 is the Maurer-Cartan form of $PL(m)$, we have $d\Lambda_1 + \frac{1}{2}[\Lambda_1, \Lambda_1] = 0$. Hence (Q, ω) is flat if and only if $(Q \times PL(m), \omega \times \Lambda_1)$ is flat. This argument also proves the equivalence of flatness between $(Q, *\omega)$ and $(Q \times PL(n), *\omega \times \Lambda_1)$.

Finally we consider the base space. Now assume (1). We fix the complementary subspace $\mathfrak{m}$ of $\mathfrak{g}_{<v>}$ in $\mathfrak{g}$, then the natural inclusion $F : G \to PL(l)$ gives the linear isomorphism $\hat{dF} : \mathfrak{m} \to M(n, m)$. We denote the isotropy representation of $G/G_{<v>}$ by $\rho : G_{<v>} \hookrightarrow GL(\mathfrak{m})$. Then $(Q/ker\rho, \theta)$ gives $\rho(G_{<v>})$-structure $\widetilde{Q} \subset \mathcal{L}(M)$. From Proposition 2.1 we obtain the induced Grassmannian Cartan connection (Q', ω'), which induces a gr-structure $(P_tM, [\chi])$ on M. Here P_tM is a $GL(n) \otimes GL(m)$-structure on M. Now by proposition 2.5 the natural inclusion $\iota : Q \to Q'$ induces the injective $\bar{\iota} : \widetilde{Q} \to P_tM$. Consequently we obtain the following diagram.

$$\begin{tikzcd}
Q \arrow[r, hook, "i"'] \arrow[d, "\rho"'] \arrow[dr, phantom, "\circlearrowleft"] & Q' \arrow[d, "\tilde{\rho}"'] \\
\widetilde{Q} \arrow[r, hook, "\bar{\iota}"'] & P_tM.
\end{tikzcd}$$

Thus we obtain the map $\bar{\iota} \circ \rho : Q \to P_tM$ corresponding to the restriction $\tilde{\rho}|_{G_{<v>}}$. For a matrix $A \in GL(m)$ denote by $\bar{A}$ the image of the homomorphism $GL(m) \to PL(m)$. We define the map $\Phi : Q \times PL(m)/G \times PL(m)_v \to P_tM/GL(n) \otimes GL(1)$ by

$$\Phi : (u, \bar{A})G \times PL(m)_v \mapsto \bar{\iota} \circ \rho(u) I_n \otimes A^{-1} GL(n) \otimes GL(1).$$

This definition is well defined. Moreover the map Φ is a diffeomorphism. Now we observe that $PL(m)$ naturally acts on $Q \times PL(m)/G \times PL(m)_v$ and $P_tM/GL(n) \otimes GL(1)$ on the right as follows: $(u, \bar{A})G \times PL(m)_v \cdot \bar{B} := (u, \overline{B^{-1}A})G \times PL(m)_v$ and $x\ GL(n) \otimes GL(1) \cdot \bar{B} := x\ I_n \otimes B\ GL(n) \otimes GL(1)$. We can check that by these actions both $Q \times PL(m)/G \times PL(m)_v$ and $P_tM/GL(n) \otimes GL(1)$ can be regarded as principal fiber bundles over M with structure group $PL(m)$. Then Φ gives a bundle isomorphism. If we have $m = 1$, then the base space of $(Q \times PL(m), \omega \times \Lambda_1)$ is same with M, and the base space of $(Q \times PL(n), *\omega \times \Lambda_1)$ is isomorphic to $\mathcal{L}(M)/GL(1)$. □

This proposition is described by the following commutative diagram. We assume (Q, ω) is a Grassmannian Cartan connection over M.

$$\begin{tikzcd}
& (Q \times PL(m), \omega \times \Lambda_1) \arrow[dd, "\pi_m"] \\
(Q, \omega) \arrow[ur] \arrow[dr] \arrow[d, "\rho"'] & \\
P_t M \arrow[r, "r_m"'] \arrow[d, "\pi"'] & M_m \arrow[dl, "\bar{\pi}"] \\
M & \\
(Q, *\omega) \arrow[u] &
\end{tikzcd}$$

The manifold M_m denotes a quotient manifold $P_tM/GL(n) \otimes GL(1)$, and we denote by r_m the projection $P_tM \to P_tM/GL(n) \otimes GL(1)$. The manifold M_m is naturally isomorphic with $Q \times_{PL(l)_{<v>}} PL(m)$. Thus we obtain a natural projection from Q to M_m and there exists a natural inclusion $Q \to Q \times PL(m)$. Denote by π_m a projection from $Q \times PL(m)$ to the quotient manifold $Q \times PL(m)/PL(l) \times PL(m)_v$. From the proof of Proposition 3.3 the quotient space $Q \times PL(m)/PL(l) \times PL(m)_v$ is identified with M_m by the bundle isomorphism given by $\pi_m(z, g) \leftrightarrow r_m \circ \rho(z)g^{-1}$.

Definition 3.5. *Let (Q, ω) satisfy the same assumption as Proposition 3.1. Then we call the transformation*

$$(Q \times PL(m), \omega \times \Lambda_1) \leftrightarrow (Q \times PL(n), *\omega \times \Lambda_1)$$

a castling transformation of projective structures. If $(Q \times PL(m), \omega \times \Lambda_1)$ induces a projective Cartan connection, then $(Q \times PL(n), *\omega \times \Lambda_1)$ also induces a projective Cartan connection.

We note that the model space of $(Q \times PL(m), \omega \times \Lambda_1)$ is $G \times PL(m)/G \times PL(m)_v$, and the one of $(Q \times PL(n), *\omega \times \Lambda_1)$ is $*G \times PL(n)/*G \times PL(n)_{v^\perp}$, where v is a fixed point of $V_{m,l}$ and $v^\perp$ is one of $V_{n,l}$.

4 Investigation of castling transformations

4.1 G-structures and linear connections

Now we describe the castling transformation of projective structures from the view point of G-structures and linear connections. Firstly we consider the differential geometry corresponding to the transformation $G \leftrightarrow *G$ for a Lie subgroup G of $PL(l)$. Let (Q, ω) be a Grassmannian Cartan connection of type (n, m) over M. By Proposition 3.3 $(Q, *\omega)$ gives a Grassmannian Cartan connection of type (m, n) over M.

Proposition 4.1. *The $GL(n) \otimes GL(m)$-structure on M induced from (Q, ω) is naturally identified with the $GL(m) \otimes GL(n)$-structure on M induced from $(Q, *\omega)$.*

Proof. By the isomorphism $* : PL(l) \to PL(l)$ the model space $PL(l)/PL(l)_{<v>}$ is a subgeometry of $PL(l)/PL(l)_{<v^\perp>}$ and the Cartan connection (Q, ω) is a subgeometry of $(Q, *\omega)$. The Lie algebra $\mathfrak{sl}(l)$ is decomposed into $M(n,m) \oplus \mathfrak{sl}(l)_{<v>}$ and $M(m,n) \oplus \mathfrak{sl}(l)_{<v^\perp>}$. The isomorphism $*$ of $\mathfrak{sl}(l)$ induces a linear isomorphism $M(n,m) \to M(m,n)$ defined by $X \mapsto -{}^tX$. Since $M(n,m)$(resp. $M(m,n)$) is naturally identified with $\boldsymbol{R}^n \otimes \boldsymbol{R}^m$ (resp. $\boldsymbol{R}^m \otimes \boldsymbol{R}^n$), $*$ induces the linear isomorphism $*$: $\boldsymbol{R}^n \otimes \boldsymbol{R}^m \to \boldsymbol{R}^m \otimes \boldsymbol{R}^n$ defined by $\xi \otimes \eta \mapsto -\eta \otimes \xi$. Let P_tM be a $GL(n) \otimes GL(m)$-structure induced from (Q, ω) and P_tM' be a $GL(m) \otimes GL(n)$-structure induced from $(Q, *\omega)$. Then by Proposition 2.5 there is a natural bundle isomorphism $t : P_tM \to P_tM'$ defined by $t : x \mapsto x \circ *$, where x is regarded as a linear isomorphism $\boldsymbol{R}^n \otimes \boldsymbol{R}^m \to T_pM$. ☐

Now we describe a projective structure obtained by a castling transformation from the view point of linear connections. Let P_tM be a Grassmannian structure. Then there exists a local section $\sigma : U \hookrightarrow P_tM$ for an open subset U of M. Since $\sigma(p)$ gives a linear isomorphism from $\boldsymbol{R}^n \otimes \boldsymbol{R}^m$ to T_pM, we obtain a vector field X_{α_i} defined by $X_{\alpha_i p} := \sigma(p)(e_\alpha \otimes f_i)$. There exists a unique admissible class of linear connections in P_tM and we choose one representative χ of that class. When we denote by θ the canonical form of P_tM, we have $\theta(\sigma_* X_{\alpha_i}) = e_\alpha \otimes f_i$. Define the Christoffel's symbols $\{\Gamma^{\gamma_k}_{\alpha_i\beta_j}\}$ corresponding to χ by

$$\nabla_{X_{\alpha_i}} X_{\beta_j} = \sum_{\gamma_k} \Gamma^{\gamma_k}_{\alpha_i\beta_j} X_{\gamma_k}.$$

Let π be the projection $P_tM \to M$ and we denote by $\bar{\pi}$ the induced projection $M_m := P_tM/GL(n) \otimes GL(1) \to M$. Denote by r_m a projection from P_tM to M_m. The connection Γ corresponding to χ in P_tM induces a connection Γ_m in M_m defined by $\Gamma_m(r_m(x)) := r_{m*}(\Gamma(x))$. For a matrix $A \in \mathfrak{gl}(m)$ we denote by $\bar{A}$ the image of the linear map $\mathfrak{gl}(m) \to \mathfrak{sl}(m)$. Since the value of the connection form $\chi(Y)$ is written as $\chi_n(Y) \otimes I_m + I_n \otimes \chi_m(Y)$ for $Y \in T_xP_tM$, the connection form χ' corresponding to Γ_m is given by $\chi'(r_{m*}Y) := \overline{\chi}_m(Y)$. By the horizontal lift of X_{α_i} with respect to χ_m we obtain a vector field in $\overline{\pi}^{-1}(U)$ and denote it by $X^*_{\alpha_i}$. On the other hand about the vertical direction, for the element $Y \in \mathfrak{sl}(m)$ we define the vector field Y on $\bar{\pi}^{-1}(U)$ by

$$Y_{r_m \circ \sigma(p) g} := R_{g*}(Y^*_{r_m \circ \sigma(p)}).$$

Let (Q, ω) be a normal Grassmannian Cartan connection constructed from (P_tM, χ). Then in Proposition 3.3 we showed that $(Q \times PL(m),\ \omega \times \Lambda_1)$ is a Cartan connection of type $PL(l) \times PL(m)/PL(l) \times PL(m)_v$, which is a subgeometry of the projective space $PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)/PL(\boldsymbol{R}^l \times \boldsymbol{R}^m)_v$. By Proposition 2.1 we obtain a projective Cartan connection $(\widetilde{Q}_m, \widetilde{\omega}_m)$ induced from $(Q \times PL(m),\ \omega \times \Lambda_1)$. Then $(\widetilde{Q}_m, \widetilde{\omega}_m)$ induces a linear connection $\widetilde{\chi}$ on M_m by using an injective homomorphism $h_m : \mathcal{L}(M_m) \to \widetilde{Q}_m$.

Proposition 4.2. *Let $[\widetilde{\chi}]$ be the projective structure on M_m induced from the Cartan connection $(Q \times PL(m),\ \omega \times \Lambda_1)$. Let $\widetilde{\nabla}$ be the covariant derivative corresponding to $\widetilde{\chi}$. With respect to* *the basis of vector fields* $\{X^*_{\alpha_i}, Y\}$ *on* $\bar{\pi}^{-1}(U)$, $\widetilde{\nabla}$ *is described as follows:*

$$\begin{aligned}
\widetilde{\nabla}_{X^*_{\alpha_i}} X^*_{\beta_j} &= \sum_{\gamma_k} \Gamma^{\ \gamma_k}_{\alpha_i\beta_j} X^*_{\gamma_k} - {}^t\overline{\begin{pmatrix} & J_{\sigma(p)}(e_{\alpha_i})_{\beta_1} & \\ 0\cdots 0 & \vdots & 0\cdots 0 \\ & J_{\sigma(p)}(e_{\alpha_i})_{\beta_m} & \end{pmatrix}}, \\
\widetilde{\nabla}_{X^*_{\alpha_i}} Y &= -\sum_{k=1}^{m} Y^k_i X^*_{\alpha_k} + [\chi_m(\sigma_* X_{\alpha_i}), Y], \\
\widetilde{\nabla}_Y X^*_{\beta_j} &= -\sum_k Y^k_j X^*_{\beta_k}, \\
\widetilde{\nabla}_Y Z &= -\overline{YZ}.
\end{aligned}$$

We note that about the above $m \times m$ matrix its component is zero without the j-th row, and J is a tensor field introduced in section 2.1, which is defined to be

$$J_{\sigma(p)}(e_{\alpha_i})_{\beta_k} = \frac{1}{(l^2-4)l}\{(2-l^2)\mathrm{Ric}(e_{\alpha_i}, e_{\beta_k}) - l\mathrm{Ric}(e_{\beta_i}, e_{\alpha_k}) - l\mathrm{Ric}(e_{\alpha_k}, e_{\beta_i}) - 2\mathrm{Ric}(e_{\beta_k}, e_{\alpha_i})\},$$

where Ric is the Ricci tensor field of χ.

REMARK 4.3. We assumed that a (Q, ω) is a normal Grassmannian Cartan connection of type (n, m), where $n \geq 2$ and $m \geq 2$. If (Q, ω) is a projective Cartan connection, then a Cartan connection $(Q \times PL(n), *\omega \times \Lambda_1)$ induces a linear connection $\widetilde{\chi}$ on M_n. Then we can describe $\widetilde{\chi}$ in the same way.

Proof. We recall that the base space M_m denotes $P_tM/GL(n) \otimes GL(1)$, and it is isomorphic to $Q \times PL(m)/PL(l) \times PL(m)_v$ as a $PL(m)$-bundle. We decompose the Lie algebra $\mathfrak{sl}(l) \times \mathfrak{sl}(m)$ into $(M(n,m) \times \mathfrak{sl}(m)) \oplus (\mathfrak{sl}(l) \times \mathfrak{sl}(m)_v)$. We denote by ρ_c the linear isotropy representation of $PL(l) \times PL(m)_v$ on the tangent space $\mathfrak{sl}(l) \times \mathfrak{sl}(m)/\mathfrak{sl}(l) \times \mathfrak{sl}(m)_v \cong M(n,m) \times \mathfrak{sl}(m)$. Thus we obtain the map $\rho_c : PL(l) \times PL(m)_v \to GL(M(n,m) \times \mathfrak{sl}(m))$. Then we obtain $\rho_c(PL(l) \times PL(m)_v)$-structure $(Q \times PL(m)/ker\rho_c, \theta_m)$ over M_m. We denote the projection $Q \times PL(m) \to Q \times PL(m)/ker\rho_c \subset \mathcal{L}(M_m)$ also by ρ_c. The canonical form θ_m is determined by ${\rho_c}^*\theta_m = \omega \times {\Lambda_1}_{M(n,m)\times\mathfrak{sl}(m)}$.

Since the bundle Q is the extended bundle of P_tM, we have the natural injective homomorphism $h : P_tM \hookrightarrow Q$ corresponding to $\iota : GL(n) \otimes GL(m) \hookrightarrow PL(l)_{<v>}$. Since M_m is isomorphic to $Q \times PL(m)/G \times PL(m)_v$, the bundle $h(P_tM) \times PL(m)$ gives a reduction of the structure group $PL(l) \times PL(m)_v$ of $Q \times PL(m)$ to $G_0 \times PL(m)_v$. Now we consider the bundle homomorphism $h(P_tM) \times PL(m) \to \mathcal{L}(M_m)$ obtained by the restriction of ρ_c. This bundle homomorphism is corresponding to the restriction $\rho_c : G_0 \times PL(m)_v \to GL(M(n,m) \times \mathfrak{sl}(m))$ given by

$$(\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}, *\overline{A}) \mapsto (B \otimes *A, Ad(*A))$$

Thus the restriction $\rho_c : h(P_tM) \times PL(m) \to \mathcal{L}(M_m)$ is injective. Hence $h(P_tM) \times PL(m)$ itself gives a $GL(n) \otimes GL(m)$-structure over M_m.

Now we consider the subgeometry $PL(l) \times PL(m)/PL(l) \times PL(m)_v$ of $PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)/PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)_v$. This subgeometry is given by the injective homomorphism $F : PL(l) \times PL(\boldsymbol{R}^m) \hookrightarrow PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)$ defined by $F : (x, y) \mapsto x \otimes y$. Since $\boldsymbol{R}^l \otimes \boldsymbol{R}^m$ is decomposed into $<(\frac{I_m}{0})>$

$\oplus(\frac{\mathfrak{sl}(m)}{M(n,m)})$, we have the graded decomposition $\mathfrak{sl}(\boldsymbol{R}^l \otimes \boldsymbol{R}^m) = (\frac{\mathfrak{sl}(m)}{M(n,m)}) \oplus \mathfrak{gl}(\frac{\mathfrak{sl}(m)}{M(n,m)}) \oplus (\frac{\mathfrak{sl}(m)}{M(n,m)})^*$. On the other hand we have the decomposition $\mathfrak{sl}(l) \times \mathfrak{sl}(m) = (M(n,m) \times \mathfrak{sl}(m)) \oplus (\mathfrak{sl}(l) \times \mathfrak{sl}(m)_v)$. Since $\rho_c : G_0 \times PL(m)_v \to GL(M(n,m) \times \mathfrak{sl}(m))$ is injective, we can define the homomorphism $\iota_m : \rho_c(G_0 \times PL(m)_v) \to PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)_v$ by $\iota_m : \rho_c(x,y) \mapsto F(x,y)$. Then by Lemma 2.4 we obtain the following commutative diagram

$$\begin{tikzcd}
G_0 \times PL(m)_v \arrow[r, "F"] \arrow[d, "\rho_c"', "\circlearrowleft"] & PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)_v \arrow[d, "\tilde{\rho}", "\circlearrowright"'] \\
\rho_c(G_0 \times PL(m)_v) \arrow[ur, "\iota_m"'] \arrow[r, hook, "\overline{F}"'] & GL(\frac{\mathfrak{sl}(m)}{M(n,m)}).
\end{tikzcd}$$

Moreover we consider a natural injection $\tilde{\iota}$: $GL(\frac{\mathfrak{sl}(m)}{M(n,m)}) \to PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)_v$. Then we have $\tilde{\iota} \circ \overline{F} = \iota_m$. Indeed we can directly check the equality as follows:

$$\tilde{\iota} \circ \overline{F}(\rho_c(\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}, *A)) \cdot (\frac{I_m}{0}) = (\frac{I_m}{0})$$

$$\begin{aligned}
\tilde{\iota} \circ \overline{F}(\rho_c(\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}, *A)) \cdot (\frac{Y}{\xi}) &= d\hat{F}(B\xi A^{-1}, *A^t Y^t A) \\
&= \begin{pmatrix} AYA^{-1} \\ B\xi A^{-1} \end{pmatrix}
\end{aligned}$$

$$\iota_m(\rho_c(\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}, *A)) = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \otimes *A$$

The natural extension of $(Q \times PL(m), \omega \times \Lambda_1)$ by $\widetilde{F}$ yields the projective Cartan connection $(\widetilde{Q}_m, \widetilde{\omega}_m)$. Define a bundle homomorphism $h_m : h(P_tM) \times PL(m) \to \widetilde{Q}_m$ by using $\widetilde{F}$ and ρ_c analogously with ι_m. Then h_m is a bundle homomorphism corresponding to $\iota_m : \rho_c(G_0 \times PL(m)_v) \to P(lm)_v$. Then by Proposition 2.5 we obtain the following commutative diagram:

$$\begin{tikzcd}
h(P_tM) \times PL(m) \arrow[r, "\widetilde{F}"] \arrow[d, "\rho_c"', "\circlearrowleft"] & \widetilde{Q}_m \arrow[d, "\tilde{\rho}", "\circlearrowright"'] \\
\rho_c(h(P_tM) \times PL(m)) \arrow[ur, "h_m"'] \arrow[r, hook, "t"'] & L(M_m).
\end{tikzcd}$$

By using t we can identify $\rho_c(h(P_tM) \times PL(m))$ with a subbundle of $\mathcal{L}(M_m)$ consisting of the linear isomorphisms $(\frac{\mathfrak{sl}(m)}{M(n,m)}) \to T_xM_m$. Thus we also identify the group $\rho_c(G_0 \times PL(m)_v)$ with a subgroup of $GL(\frac{\mathfrak{sl}(m)}{M(n,m)})$ via $\overline{F}$. By Lemma 2.4 the identification of these groups is given by the identification of the subspace $M(n,m) \times \mathfrak{sl}(m)$ with $(\frac{\mathfrak{sl}(m)}{M(n,m)})$ defined by $(\xi, Y) \leftrightarrow \begin{pmatrix} {}^tY \\ \xi \end{pmatrix}$. Under this identification h_m satisfies $\tilde{\rho} \circ h_m = id$, and the corresponding group homomorphism ι_m is identified with the restriction of $\tilde{\iota}$: $GL(\frac{\mathfrak{sl}(m)}{M(n,m)}) \to PL(\boldsymbol{R}^l \otimes \boldsymbol{R}^m)_v$.

The bundle homomorphism h_m enables us to compute a connection form induced from the Cartan connection $(\widetilde{Q}_m, \widetilde{\omega}_m)$. Indeed we extend h_m to the bundle homomorphism $h'_m : \mathcal{L}(M_m) \to \widetilde{Q}_m$. We denote the $\mathfrak{gl}(\frac{\mathfrak{sl}(m)}{M(n,m)})$-component of $\widetilde{\omega}_m$ by $(\widetilde{\omega}_m)_0$. Then we have $\widetilde{\chi} = {h'_m}^*(\widetilde{\omega})_0$. Obviously in the subbundle $\rho_c(h(P_tM) \times PL(m))$ of $L(M_m)$, $\widetilde{\chi} = {h_m}^*(\widetilde{\omega}_m)_0$.

Now let $\widetilde{\nabla}$ be the covariant derivative corresponding to the connection form $\widetilde{\chi}$. To compute $\widetilde{\nabla}$ from $\widetilde{\chi}$, we define the section σ_m: $\bar{\pi}^{-1}(U) \hookrightarrow \mathcal{L}(M_m)$ by

$$\sigma_m : \pi_m(h(\sigma(p)), g) \mapsto \rho_c(h(\sigma(p)), g).$$

We compute the linear frame $\rho_c(h(\sigma(p)), g^{-1})$.

By using the section $\sigma : U \to P_tM$ and the vector fields X_{α_i} on the neighborhood U we obtain the tangent vector $\sigma_* X_{\alpha_i}$ on P_tM. Let $Y \in \mathfrak{sl}(m)$ be a left invariant vector field. Then $(h_*\sigma_*X_{\alpha_i}, \overline{\chi_m(\sigma_*X_{\alpha_i})}_g)$ and $(0, Y_g)$ give tangent vectors at $(h(\sigma(p)), g)$ in $h(P_tM) \times PL(m)$. We compute the canonical form θ_m at $s = \rho_c(h(\sigma(p)), g^{-1})$.

$$\begin{aligned}
& \theta_m(\rho_{c*}(h_*(\sigma_*X_{\alpha_i}), \overline{\chi_m(\sigma_*X_{\alpha_i})}_{g^{-1}}) \\
= \; & \omega \times \Lambda_{1\,M(n,m)\times\mathfrak{sl}(m)}(h_*(\sigma_*X_{\alpha_i}), \overline{\chi_m(\sigma_*X_{\alpha_i})}_{g^{-1}}) \\
= \; & (\begin{pmatrix} -{}^t\chi_m(\sigma_*X_{\alpha_i}) & J_x(\theta(\sigma_*X_{\alpha_i})) \\ \theta(\sigma_*X_{\alpha_i}) & \chi_n(\sigma_*X_{\alpha_i}) \end{pmatrix}, \overline{\chi_m(\sigma_*X_{\alpha_i})})_{M(n,m)\times\mathfrak{sl}(m)} \\
= \; & (\theta_{\sigma(p)}(\sigma_*X_{\alpha_i}), 0) \\
= \; & (e_\alpha \otimes f_i, 0),
\end{aligned}$$

$$\theta_m(\rho_{c*}(0, Y_{g^{-1}})) = (0, Y).$$

It follows that the linear frame s of M_m is the linear isomorphism $s : M(n,m) \times \mathfrak{sl}(m) \to T_{\pi_m(h(\sigma(p), g^{-1}))}M_m$ defined by

$$\begin{aligned}
(e_{\alpha_i}, 0) \;& \mapsto \; \pi_{m*}(h_*(\sigma_*X_{\alpha_i}), \overline{\chi_m(\sigma_*X_{\alpha_i})}_{g^{-1}}) \\
(0, Y) \;& \mapsto \; \pi_{m*}(0, Y_{g^{-1}}).
\end{aligned}$$

Since

$$\pi_m(h(x), g^{-1}) = r_m(x)g,$$

the tangent vector $\pi_{m*}(h_*(\sigma_*X_{\alpha_i}), \overline{\chi_m(X_{\alpha_i})}_{g^{-1}})$ at $\pi_m(h(\sigma(p)), g^{-1})$ is equal to

$$R_{g*} r_{m*}(\sigma_* X_{\alpha_i\,\sigma(p)}) - R_{g*}(\overline{\chi_m(\sigma_*X_{\alpha_i})})^*_{q_m\circ\sigma(p)}.$$

This vector is included in the kernel of the connection form χ' on M_m. Hence we have

$$\pi_{m*}(h_*X_{\alpha_i\,\sigma(p)}, \overline{\chi_m(X_{\alpha_i})}_{g^{-1}}) = X^*_{\alpha_i\,r_m\circ\sigma(p)g}.$$

Likewise the tangent vector $\pi_{m*}(0, -Y_{g^{-1}})$ at $\pi_m(h(\sigma(p)), g^{-1})$ is equal to $R_{g*}(Y^*_{r_m\circ\sigma(p)})$, thus we have

$$\pi_{m*}(0, -Y_{g^{-1}}) = Y_{q_m(h(\sigma(p)))g}.$$

Moreover we have

$$\sigma_{m*} \circ \pi_{m*}(h_*(\sigma_* X_{\alpha_i}), \overline{\chi_m(\sigma_* X_{\alpha_i})}_{g^{-1}}) = \rho_{c*}(h_*(\sigma_* X_{\alpha_i}), \overline{\chi_m(\sigma_* X_{\alpha_i})}_{g^{-1}}),$$

$$\sigma_{m*}(\pi_{m*}(0, -Y_{g^{-1}})) = \rho_{c*}(0, -Y_{g^{-1}}).$$

Here we state one general assertion: let $\{Z_1, \dots, Z_n\}$ be a basis of vector fields on a manifold N^n, and χ be a connection form on $\mathcal{L}(N)$. We denote by θ a canonical form of $\mathcal{L}(N)$. For some open neighborhood U of N we consider a section $\sigma' : U \hookrightarrow \mathcal{L}(N)$ defined by $\sigma'(p) = (Z_{1p}, \dots, Z_{np})$. We define the Christoffel's symbols $\{\Gamma_i{}^k{}_j\}$ by

$$\sigma'^* \chi(Z_i) \theta({\sigma'}_* Z_j) = \sum_{j,k} \Gamma_i{}^k{}_j \theta({\sigma'}_* Z_k)$$

with respect to the basis $\{Z_1, \dots, Z_n\}$. Denote the covariant derivative ∇ corresponding to χ. Then we have

$$\nabla_{Z_i} Z_j := \sum_k \Gamma_i{}^k{}_j Z_k.$$

This assertion can be proved by using the fundamental correspondence between a connection form and the corresponding Christoffel's symbols with respect to the natural vector fields induced by coordinates of N.

Now we compute the covariant derivative $\widetilde{\nabla}$ on M_m. The equalities

$$\sigma_{m*} \circ \pi_{m*}(h_*(\sigma_* X_{\alpha_i}), \overline{\chi_m(\sigma_* X_{\alpha_i})}_{g^{-1}}) = \rho_{c*}(h_* X_{\alpha_i}, \overline{\chi_m(X_{\alpha_i})}_{g^{-1}})$$

and

$$\begin{aligned} & \widetilde{\chi}_m(\rho_{c*}(h_*(\sigma_* X_{\alpha_i}), \overline{\chi_m(\sigma_* X_{\alpha_i})}_{g^{-1}})) \\ = \quad & dF(\begin{pmatrix} -{}^t\chi_m(\sigma_* X_{\alpha_i}) & J_x(\theta(\sigma_* X_{\alpha_i})) \\ \theta(\sigma_* X_{\alpha_i}) & \chi_n(\sigma_* X_{\alpha_i}) \end{pmatrix}, \overline{\chi_m(\sigma_* X_{\alpha_i})})_{\mathfrak{gl}(\frac{\mathfrak{sl}(m)}{M(n,m)})}, \end{aligned}$$

yields

$$\begin{aligned} & \widetilde{\chi}_m(\sigma_{m*} \circ \pi_{m*}(h_*(\sigma_* X_{\alpha_i}), \overline{\chi_m(\sigma_* X_{\alpha_i})}_{g^{-1}})) \cdot (e_{\beta_j}, 0) \\ = \quad & (\chi_n(X_{\alpha_i}) \cdot e_{\beta_j} + e_{\beta_j} \cdot {}^t\overline{\chi_m(\sigma_* X_{\alpha_i})} + \frac{\mathrm{tr}\chi_m(\sigma_* X_{\alpha_i})}{m} e_{\beta_j},\ {}^t(\overline{J_{\sigma(p)}(e_{\alpha_i}) \cdot e_{\beta_j}}). \end{aligned}$$

We denote the matrix components of χ_m and χ_n by

$$\begin{aligned} \chi_n(\sigma_* X_{\alpha_i}) e_\beta &= \sum_\gamma \chi_n(\sigma_* X_{\alpha_i})^\gamma_\beta e_\gamma, \\ \chi_m(\sigma_* X_{\alpha_i}) f_j &= \sum_k \chi_m(\sigma_* X_{\alpha_i})^k_j f_k. \end{aligned}$$

Then from the equality

$$\chi(\sigma_* X_{\alpha_i}) e_\beta \otimes f_j = \sum_{\gamma_k} (\chi_n(\sigma_* X_{\alpha_i})^\gamma_\beta \delta^k_j + \chi_m(\sigma_* X_{\alpha_i})^k_j \delta^\gamma_\beta) e_\gamma \otimes f_k$$

The $M(n,m)$-component $\chi_n(X_{\alpha_i}) \cdot e_{\beta_j} + e_{\beta_j} \cdot {}^t\overline{\chi_m(X_{\alpha_i})} + \frac{\mathrm{tr}\chi_m(X_{\alpha_i})}{m} e_{\beta_j}$ is equal to

$$\sum_{\gamma_k} \{\chi_n(\sigma_* X_{\alpha_i})^\gamma_\beta \delta^k_j + \chi_m(\sigma_* X_{\alpha_i})^k_j \delta^\gamma_\beta\} e_{\gamma_k}.$$

The $\mathfrak{sl}(m)$-component is equal to

$$A := {}^t\overline{\begin{pmatrix} & J_{\sigma(p)}(e_{\alpha_i})_{\beta_1} & \\ 0\cdots 0 & \vdots & 0\cdots 0 \\ & J_{\sigma(p)}(e_{\alpha_i})_{\beta_m} & \end{pmatrix}}.$$

Moreover by the formula

$$\Gamma_{\alpha_i}{}^{\gamma_k}{}_{\beta_j} = \chi_n(\sigma_* X_{\alpha_i})^\gamma_\beta \delta^k_j + \chi_m(\sigma_* X_{\alpha_i})^k_j \delta^\gamma_\beta$$

we obtain

$$\widetilde{\chi}_m(\sigma_{m*} \circ \pi_{m*}(h_* \sigma_* X_{\alpha_i}, \overline{\chi_m(\sigma_* X_{\alpha_i})}_{g^{-1}})) \cdot (e_{\beta_j}, 0) = (\sum_{\gamma_k} \Gamma_{\alpha_i}{}^{\gamma_k}{}_{\beta_j} e_{\gamma_k},\ A).$$

Likewise we obtain

$$\begin{aligned} \widetilde{\chi}_m(\sigma_{m*} \circ \pi_{m*}(h_* \sigma_* X_{\alpha_i}, \overline{\chi_m(\sigma_* X_{\alpha_i})}_{g^{-1}})) \cdot (0, -Y) &= (-\sum_k Y^k_i e_{\alpha_k},\ -[\chi_m(\sigma_* X_{\alpha_i}), Y]), \\ \widetilde{\chi}_m(\sigma_{m*} \circ \pi_{m*}(0, -Y_{g^{-1}})) \cdot (e_{\beta_j}, 0) &= (-\sum_k Y^k_j e_{\beta_k},\ 0), \\ \widetilde{\chi}_m(\sigma_{m*} \circ \pi_{m*}(0, -Y_{g^{-1}})) \cdot (0, -Z) &= (0,\ \overline{YZ}). \end{aligned}$$

This proves the Proposition 4.2. □

REMARK 4.4. If the Ricci tensor of χ in P_tM is equal to 0, the geometrical meaning of the linear connection $\widetilde{\nabla}$ is clear in the sense of parallel transport of tangent vectors of M_m. However for the author it is difficult to understand the meaning of the matrix term appeared in $\widetilde{\nabla}_{X^*_{\alpha_i}} X^*_{\beta_j}$.

4.2 Normality and torsion of castling transformations

Suppose that χ is a linear connection in P_tM, (Q, ω) is a Grassmannian Cartan connection constructed from (P_tM, χ) by using an admissible homomorphism $h : P_tM \to Q$. Let Ω be the curvature form of (Q, ω). We denote the $M(n,m)$-component of Ω by S, and the $\mathfrak{gl}(n) \otimes I + I \otimes \mathfrak{gl}(m)$-component and the $M(m,n)$-component of Ω by K^0 and K^1 respectively. For $\xi, \eta \in M(n,m)$ and $z \in Q$ we express $\Omega(\check{\omega}_z(\xi), \check{\omega}_z(\eta))$ as $\Omega_z(\xi, \eta)$. Then by [Tan2, Lemma 6.1 and Proposition 7.5] the torsion tensor T of χ is computed by

$$T_x(\xi, \eta) = S_{h(x)}(\xi, \eta).$$

Let $\widetilde{\Omega}$ be the curvature form of the projective Cartan connection $(\tilde{Q}, \tilde{\omega})$. We denote the $(\frac{\mathfrak{sl}(m)}{M(n,m)})$-component of $\widetilde{\Omega}$ by $\tilde{S}$ and the torsion tensor of $\widetilde{\chi}$ by $\widetilde{T}$. Then $\widetilde{T}$ is computed by

$$\widetilde{T}_{\rho_c(h(\sigma(p)), g^{-1})}((\xi, 0), \eta, 0)) = \widetilde{S}_{\tilde{F}(h(\sigma(p)), g^{-1})}(\begin{pmatrix} 0 \\ \hline \xi \end{pmatrix}, \begin{pmatrix} 0 \\ \hline \eta \end{pmatrix}).$$

Proposition 4.5. *The projective Cartan connection $(\widetilde{Q}, \widetilde{\omega})$ obtained by the castling transformation is generally not normal. Indeed the torsion tensor field $\widetilde{T}$ of $\widetilde{\chi}$ is given as follows:*

$$\widetilde{T}_{\rho_c(h(\sigma(p)),g^{-1})}((\xi,0),(\eta,0)) = (T_{\sigma(p)}(\xi,\eta), {}^t\overline{K^0_{m\,h(\sigma(p))}(\check{\omega}(\xi),\check{\omega}(\eta))} \qquad (\xi,\eta \in M(n,m)).$$

However the castling transformation is a transformation of the projective structures not only of linear connections.

Proof. For elements ξ, η of $M(n,m)$ the curvature form $\widetilde{\Omega}$ is computed as follows:

$$\begin{aligned}\widetilde{\Omega}(\tilde{\check{\omega}}\begin{pmatrix} 0 \\ \hline \xi \end{pmatrix}, \tilde{\check{\omega}}\begin{pmatrix} 0 \\ \hline \eta \end{pmatrix}) &= -\widetilde{\omega}[\tilde{\check{\omega}}\begin{pmatrix} 0 \\ \hline \xi \end{pmatrix}, \tilde{\check{\omega}}\begin{pmatrix} 0 \\ \hline \eta \end{pmatrix}] \\ &= -dF(\omega[\check{\omega}(\xi), \check{\omega}(\eta)], 0) \\ &= dF(\begin{pmatrix} K^0_{m\,h(x)}(\xi,\eta) & K^1{}_{h(x)}(\xi,\eta) \\ S_{h(x)}(\xi,\eta) & K^0_{n\,h(x)}(\xi,\eta) \end{pmatrix}, 0).\end{aligned}$$

Thus we have

$$\widetilde{S}_{\tilde{F}(h(x),g^{-1})}(\begin{pmatrix} 0 \\ \hline \xi \end{pmatrix}, \begin{pmatrix} 0 \\ \hline \eta \end{pmatrix}) = \begin{pmatrix} \overline{K^0_{m\,h(x)}(\xi,\eta)} \\ \hline S_{h(x)}(\xi,\eta) \end{pmatrix}.$$

Hence generally the projective Cartan connection $(\widetilde{Q}, \widetilde{\omega})$ is not normal. Now let χ' be a linear connection belonging to the admissible class $[\chi]$. The connection χ' gives rise to a normal Cartan connection (Q, ω') which is isomorphic to (Q, ω). The Cartan connections $(Q \times PL(m), \omega \times \Lambda_1)$ and $(Q \times PL(m), \omega' \times \Lambda_1)$ are isomorphic, hence the extended Cartan connections $(\widetilde{Q}, \widetilde{\omega})$ and $(\widetilde{Q}, \widetilde{\omega'})$ are also isomorphic. The Cartan connection $(\widetilde{Q}, \widetilde{\omega})$ is not normal generally as we have already investigated it by the torsion. However since $(\widetilde{Q}, \widetilde{\omega})$ and $(\widetilde{Q}, \widetilde{\omega'})$ are isomorphic, from the proof of [Tan2, Theorem 9.2] the induced linear connections $\widetilde{\chi}$ and $\widetilde{\chi}'$ by admissible homomorphisms $\mathcal{L}(M_m) \to \widetilde{Q}$ are projectively equivalent. These discussion proves the proposition. □

Conventionally a projective structure is a projective equivalence class of torsion-free linear connections. On the other hand a linear connection $\widetilde{\chi}$ on M_m is not torsion-free generally. However according to Proposition 4.5 a gr-structure $(P_tM, [\chi])$ gives rise to a geometric structure $[\widetilde{\chi}]$ naturally (see §2.1 for the projective equivalence relation). Thus we call our transformation a castling transformation of projective structures.

Now we consider a projective structure $(\mathcal{L}(M), [\chi])$ on n-dimensional manifold M. Then we can construct a normal projective Cartan connection (Q, ω) of type $PL(n+1)/PL(n+1)_v$, where v is an element of the projective space $P(\boldsymbol{R}^{n+1})$ being equivalent to $Gr_{1,n+1}$. Thus by the castling transformation $(Q, \omega) \to (Q \times PL(n), *\omega \times \Lambda_1)$ we obtain a projective Cartan connection $(\widetilde{Q}, \widetilde{*\omega})$ over M_n and it induces a projective structure $[\widetilde{\chi}]$ on M_n. Then we have the following.

Corollary 4.6. *The torsion of $[\widetilde{\chi}]$ is given by the Weyl's projective curvature tensor of $[\chi]$.*

Proof. Let $h : \mathcal{L}(M) \to Q$ be the natural inclusion. From the proof of Proposition 4.2 $h(\mathcal{L}(M)) \times PL(n)$ is regarded as a subbundle of $L(M_n)$ by the canonical form $\theta_n := *\omega \times \Lambda_1{}_{M(1,n)\times\mathfrak{sl}(n)}$. Thus we obtain the inclusion $\rho_c : h(\mathcal{L}(M)) \times PL(n) \to L(M_n)$. We denote by $\widetilde{S}$ the $(\frac{M(1,n)}{\mathfrak{sl}(n)})$-component of the curvature form $\widetilde{\Omega}$ of $(\widetilde{Q}, \widetilde{*\omega})$. From the proof of Proposition 4.5 the value of

$\widetilde{\Omega}$ at $\widetilde{F}(h(x), g^{-1})$ is computed as follows

$$\begin{aligned}\widetilde{\Omega}(\begin{pmatrix}\frac{\xi}{0}\end{pmatrix}, \begin{pmatrix}\frac{\eta}{0}\end{pmatrix}) &= -dF(*\omega[*\check{\omega}(\xi), *\check{\omega}(\eta)], 0) \\ &= dF(- * \omega[\phi_*\check{\omega}(-{}^t\xi), \phi_*\check{\omega}(-{}^t\eta)], 0) \\ &= dF({}^t\omega[\check{\omega}(-{}^t\xi), \check{\omega}(-{}^t\eta)], 0) \\ &= dF(-{}^t\begin{pmatrix} K^0_{1\,h(x)}({}^t\xi, {}^t\eta) & K^1{}_{h(x)}({}^t\xi, {}^t\eta) \\ S_{h(x)}({}^t\xi, {}^t\eta) & K^0_{n\,h(x)}({}^t\xi, {}^t\eta)\end{pmatrix}, 0).\end{aligned}$$

Here we used the equality $*\check{\omega}(\xi) = \phi_*\check{\omega}(-{}^t\xi)$. In the above $K^0_n - K^0_1 I_n$ is the $\mathfrak{gl}(n)$-component of the curvature form of (Q, ω) and S is a Torsion tensor of χ, which is assumed to be zero. Then $K^0_n - K^0_1 I_n$ is shown to coincide with the Weyl's projective curvature tensor field W of type $\begin{pmatrix}1\\3\end{pmatrix}$ on M (see [Tan1, p.10]). We recall the definition of W in the proof of Proposition 5.2. Since $K^0_1 = -\mathrm{tr}K^0_n$ and $\mathrm{tr}W({}^t\xi, {}^t\eta) = 0$, we have $K^0_n = W$ and $K^0_1 = 0$. Hence we have

$$\widetilde{T}_{\rho_c(\phi(h(x)), g^{-1})}((\xi, 0), (\eta, 0)) = (0, -W_x({}^t\xi, {}^t\eta)) \quad (\xi, \eta \in \mathfrak{gl}(1, n)).$$

□

5 Product connections

Let $(P_tM, [\chi])$ be a gr-structure and (Q, ω) be a normal Grassmannian Cartan connection constructed from $(P_tM, [\chi])$. In Proposition 4.2 we described a linear connection $\widetilde{\nabla}$ on M_m induced by $(Q \times PL(m), \omega \times \Lambda_1)$. Now with respect to the connection $\widetilde{\nabla}$ we investigate the fibers of M_m.

Proposition 5.1. *Concerning the projective structure* $[\widetilde{\nabla}]$ *on* M_m *each fiber of* M_m *is auto-parallel submanifold and the induced projective structure coincides with an invariant flat projective structure on* $PL(m)$.

Proof. Let $\bar{\pi} : M_m \to M$ be the projection. From the description of $\widetilde{\nabla}$ in Proposition 4.2 we directly see that for each point p of M the fiber $\bar{\pi}^{-1}(p)$ is auto-parallel. Thus a linear connection is induced on $\bar{\pi}^{-1}(p)$ which is given as follows: $\widetilde{\nabla}_Y Z = -\overline{YZ}$ where Y and Z are vector fields on $\bar{\pi}^{-1}(p)$ defined in 4.2. We consider the diffeomorphism $\phi : \bar{\pi}^{-1}(p) \to PL(m)$ defined by

$$\phi : r_m \circ \sigma(p)g \mapsto g^{-1}.$$

Then about the differential ϕ_*, for $Y \in \mathfrak{sl}(m)$ we have

$$\phi_*(R_{g*}(Y)^*_{r_m\circ\sigma(p)}) = -Y_{g^{-1}}.$$

Hence the vector field Y on $\bar{\pi}^{-1}(U)$ is mapped into the left invariant vector field $-Y$. Pulling back the connection $\widetilde{\nabla}$ by ϕ^{-1} yields the linear connection ∇_m on $PL(m)$, which is left invariant and defined by

$$\nabla_{mY}Z = \overline{YZ} \quad (Y, Z \in \mathfrak{sl}(m)).$$

The strait forward computation of the torsion and the the Weyl's projective curvature tensor field show that the connection ∇_m is torsion free and projectively flat. This connection coincides with the left invariant flat projective structure on $PL(m)$ constructed in [Aga]. □

Thus by Proposition 4.2 and 5.1 we see that the connection $\widetilde{\nabla}$ on M_m consists of a linear connection χ in P_tM and the invariant flat projective structure ∇_m on $PL(m)$. As the base space M_m is locally trivial, it is natural to ask whether the product connection of χ and ∇_m also gives a projectively flat connection.

Proposition 5.2. *The product connection of χ and ∇_m are not projectively flat. More generally let (M, ∇) be an affinely flat manifold and (M', ∇') be a projectively flat manifold such that its Ricci tensor field does not vanish. Then the product connection $\nabla \times \nabla'$ on $M \times M'$ is not projectively flat.*

Proof. We denote the torsion tensor, curvature tensor and Ricci tensor of ∇ (resp. ∇') respectively by T (resp. T'), R (resp. R') and Ric (resp. Ric$'$), and the ones of $\nabla \times \nabla'$ by $\overline{T}$, $\overline{R}$ and $\overline{\mathrm{Ric}}$. About these tensors we recall the basic formulas:

$$\begin{aligned}
\overline{T}((X,X'),(Y,Y')) &= (T(X,Y), T'(X',Y')) \\
\overline{R}((X,X'),(Y,Y'))(Z,Z') &= (R(X,Y)Z, R'(X',Y')Z') \\
\overline{\mathrm{Ric}}((X,X'),(Y,Y')) &= \mathrm{Ric}(X,Y) + \mathrm{Ric}'(X',Y'),
\end{aligned}$$

where $X, Y, Z \in T_pM$ and $X', Y', Z' \in T_{p'}M'$. The projective curvature tensor of Weyl W is defined by

$$W(X,Y)Z = R(X,Y)Z + [P(X,Y) - P(Y,X)]Z - [P(Y,Z)X - P(X,Z)Y],$$

where $P(X,Y) = \frac{1}{n^2-1}[n\mathrm{Ric}(X,Y) + \mathrm{Ric}(Y,X)]$ (see [NS]). About these tensors we have the following relationship:

$$\overline{P}((X,X'),(Y,Y')) = P(X,Y) + P(X',Y')$$

$$\begin{aligned}
&\overline{W}((X,X'),(Y,Y'))(Z,Z') \\
= \;& (W(X,Y)Z + [P'(X',Y') - P(Y',X')]Z - P'(Y',Z')X + P'(X',Z')Y, \\
& W'(X',Y')Z' + [P(X,Y) - P(Y,X)]Z' - [P(Y,Z)X' - P(X,Z)Y'])
\end{aligned}$$

Easily we can show that $\mathrm{Ric}(X,Y) = 0$ if and only if $P(X,Y) = 0$ and $P(X,Y) - P(Y,X) = 0$. Now we assume that ∇ is affinely flat and ∇' is projectively flat but $\mathrm{Ric} \neq 0$. Then we have

$$\overline{W}((X,X'),(Y,Y'))(Z,Z') = ([P'(X',Y') - P(Y',X')]Z - [P'(Y',Z')X - P'(X',Z')Y], 0).$$

Since the tensor Ric$'$ is not equal to zero, there exists $X', Y' \in T_{p'}M'$ such that $P'(X',Y') - P'(Y',X') \neq 0$ or $P'(X',Y') \neq 0$. We assume that $P'(X',Y') - P'(Y',X') \neq 0$. Then putting $Y = 0$ and $X = 0$ yields

$$\overline{W}((0,X'),(0,Y'))(Z,Z') = ([P'(X',Y') - P(Y',X')]Z, 0),$$

which is not equal to zero. Likewise for the case $P'(X',Y') \neq 0$ we can check $\overline{W} \neq 0$. Since ∇_m is not Ricci flat, we obtain our assertion. □

We note that under the assumption of (M', ∇') being projectively flat, ∇' is Ricci flat if and only if ∇' is affinely flat. On dimension 2 we do not need the assumption of projectively flatness.

Proposition 5.3. *Let ∇' be a linear connection on $PL(m)$ projectively equivalent to ∇_m. Then ∇' is not affinely flat.*

Proof. By a result of [Aga] a left invariant flat projective structure ($\mathcal{L}(L)$, $[\chi]$) on a Lie group L of dimension n give rise to a Lie algebra representation as follows: We fix a frame $\tilde{o}$ of tangent space T_eL and define a map $\tilde{j} : L \to \mathcal{L}(L)$ by a left translation of $\tilde{o}$. Then the Lie algebra representation $f : \mathfrak{l} \to \mathfrak{sl}(n+1)$ corresponding to χ is given by

$$f(X) = \begin{pmatrix} -\frac{\mathrm{tr}\chi(\tilde{j}_* X)}{n+1} & J(\theta(\tilde{j}_* X)) \\ \theta(\tilde{j}_* X) & \chi(\tilde{j}_* X) - \frac{\mathrm{tr}\chi(\tilde{j}_* X)}{n+1} I_n \end{pmatrix},$$

where J is a tensor field of type $(0, 2)$ defined by the formula 2.2 in the case $m = 1$. This representation satisfies the condition of (P)-homomorphism introduced in [Aga]. Conversely when a (P)-homomorphism f corresponding to a left invariant flat projective structure on L is given, there exists a left invariant projectively flat affine connection χ on L which gives rise to f. We consider the Lie algebra homomorphism $f : \mathfrak{sl}(m) \to \mathfrak{sl}(\mathfrak{gl}(m))$ defined by

$$f : X \mapsto X \otimes I_m.$$

We decompose $\mathfrak{gl}(m)$ into $< I_m > \oplus\ \mathfrak{sl}(m)$ and with respect to this decomposition $\mathfrak{sl}(\mathfrak{gl}(m))$ is decomposed into the graded Lie algebra $\mathfrak{g}_{-1}$, $\mathfrak{g}_0$, $\mathfrak{g}_1$ in a similar way to the Grassmannian manifold Gr_{1,m^2}. We regard $\mathfrak{g}_1$ and $\mathfrak{g}_{-1}$ as $\mathfrak{sl}(m)$. Moreover $\mathfrak{g}_1$ is identified with $\mathfrak{g}^*_{-1}$ by

$$<\xi, Y>:= \mathrm{tr}(\xi^t Y) \quad (\xi \in \mathfrak{g}_1, Y \in \mathfrak{g}_{-1}).$$

Then a left invariant projectively flat affine connection χ on $PL(m)$ corresponding to f is given by $\chi(\tilde{j}_* X)\theta(\tilde{j}_* Y) = f_0(X)Y = \overline{XY}$. Thus f corresponds to the invariant flat projective structure on $PL(m)$ given by ∇_m. From the assumption ∇' is projectively equivalent to ∇_m, the projective structure $[\nabla']$ is also invariant flat. Hence ∇' give rise to a Lie algebra homomorphism $f' : \mathfrak{sl}(m) \to \mathfrak{sl}(\mathfrak{gl}(m))$. Since ∇' is projectively equivalent to ∇_m, f' is projectively equivalent to f. It follows that there exists $\xi \in \mathfrak{g}_1$ such that

$$f'_1(X) = f_1(X) + [\xi, f_0(X)] + \frac{1}{2}[\xi, [\xi, f_{-1}(X)]] \quad \text{(see [Aga])}.$$

We show that f'_1 is not identically zero. From the definition of $f : \mathfrak{sl}(m) \to \mathfrak{sl}(\mathfrak{gl}(m))$ we obtain the following:

$$\begin{aligned} f_1(X)Y &= \frac{\mathrm{tr}XY}{m} \\ [\xi, f_0(X)]Y &= \mathrm{tr}({}^t\xi XY) \\ [\xi, [\xi, f_{-1}(X)]]Y &= -2\ \mathrm{tr}{}^t\xi X \mathrm{tr}{}^t\xi Y \end{aligned}$$

By using this formula we compute $f'_1(X)Y$ for the following four cases. We express ξ as the $m \times m$ matrix (a^i_j), where a^i_j denotes the (i, j)-component of ξ.

$$\begin{array}{cccc} & X & Y & f'_1(X)Y \\ (1) & E^i_i - E^m_m & E^m_i & a^i_m(1 - (a^i_i - a^m_m)) \\ (2) & E^i_i - E^m_m & E^i_m & a^m_i(1 + (a^i_i - a^m_m)) \\ (3) & E^m_i & E^i_m & \frac{1}{m} + a^i_i - a^i_m a^m_i \\ (4) & E^i_m & E^m_i & \frac{1}{m} + a^m_m - a^m_i a^i_m, \end{array}$$

where E^l_k denotes the $m \times m$ matrix B whose (p, q)-component b_{pq} is equal to $\delta^p_k \delta^q_l$, and the above index i runs from 1 to m. Now we assume that the equation $f'_1(X)Y = 0$ holds for all $X, Y \in \mathfrak{sl}(m)$. Then for a fixed i $(1 \leq i \leq m-1)$ the equation $f'_1(X)Y = 0$ of the case (1) has the two solution $a^i_m = 0$ or $a^i_i - a^m_m = 1$. If we assume $a^i_i - a^m_m = 1$, then we obtain $a^m_i = 0$ from (2), $a^i_i = -\frac{1}{m}$ from (3) and $a^m_m = -\frac{1}{m}$ from (4). However this is a contradiction. Thus we have $a^i_m = 0$ for $1 \leq i \leq m-1$. Then from (3) and (4) we obtain $a^i_i = -\frac{1}{m}$ for $1 \leq i \leq m$. This contradicts the assumption $\mathrm{tr}\xi = 0$. Therefore $f'_1(X)Y \neq 0$. From the construction of f' we have

$$f'_1(X)Y \quad = \quad J'(\theta(\tilde{j}_* X_e))(Y_e) \qquad (X, Y \in \mathfrak{sl}(m)).$$

From the equality

$$J'(\theta(\tilde{j}_* X_e))(Y_e) = -\frac{1}{n^2-1}(n\mathrm{Ric}_e(X,Y) + \mathrm{Ric}_e(Y,X)),$$

We have $f'_1(X)Y = -P'_e(X,Y)$. Since $P'(X,Y) = 0$ if and only if $\mathrm{Ric}'(X,Y) = 0$, we conclude $\mathrm{Ric}' \neq 0$. It follows that ∇' is not affinely flat. $\square$

From Propositions 5.2 and 5.3 we obtain the following Corollary.

Corollary 5.4. *Let ∇ be a affinely flat connection on M and $[\nabla_m]$ be the invariant flat projective structure on $PL(m)$ given by Proposition 5.1. For any linear connection ∇'_m projectively equivalent to ∇_m the product connection $\nabla \times \nabla'_m$ does not give a flat projective structure on $M \times PL(m)$.*

On the other hand let $(P_t M, [\chi])$ be a gr-structure of type (n, m) on M and assume the corresponding Grassmannian Cartan connection is flat. Then the linear connection $\widetilde{\nabla}$ on M_m, which is constructed from χ and ∇_m on $PL(m)$, gives a flat projective structure $[\widetilde{\nabla}]$. Especially when $m = 1$ from a flat projective structure $(\mathcal{L}(M), [\chi])$ on M, we obtain a flat projective structure $[\widetilde{\nabla}]$ on M_n again.

6 Successive castling transformations

In this section we give two fundamental procedures to do castling transformations successively. The product group $PL(l) \times \prod^j_{i=1} PL(k_i)$ naturally acts on $P(\boldsymbol{R}^l \otimes \otimes^j_{i=1} \boldsymbol{R}^{k_i})$ by the tensor product, namely via the inclusion $\imath : PL(l) \times \prod^j_{i=1} PL(k_i) \hookrightarrow PL(\boldsymbol{R}^l \otimes \otimes^j_{i=1} \boldsymbol{R}^{k_i})$ given by $\imath(\bar{g}, \bar{A}_1, \dots, \bar{A}_j) = g \otimes A_1 \otimes \cdots \otimes A_j$. We denote the natural basis of $\boldsymbol{R}^l, \boldsymbol{R}^{k_1}, \dots, \boldsymbol{R}^{k_j}$ by $\{e_{i_0}\}^l_{i_0=1}, \{e_{i_1}\}^{k_1}_{i_1=1}, \dots, \{e_{i_j}\}^{k_j}_{i_j=1}$. Then a point w in $\boldsymbol{R}^l \otimes \otimes^j_{i=1} \boldsymbol{R}^{k_i}$ is written by $w = \sum_{i_0, i_1, \dots, i_j} C_{i_0 i_1 \cdots i_j} e_{i_0} \otimes e_{i_1} \otimes \cdots \otimes e_{i_j}$ where $C_{i_0 i_1 \cdots i_j}$ is a coefficient. Now let σ be an element of symmetric group of $\{1, 2, \cdots j\}$. Then σ induces a natural linear isomorphism $\boldsymbol{R}^l \otimes \otimes^j_{i=1} \boldsymbol{R}^{k_i} \to \boldsymbol{R}^l \otimes \otimes^j_{i=1} \boldsymbol{R}^{\sigma(k_i)}$, which is defined by $\sigma(w) = \sum_{i_0, i_1, \dots, i_j} C_{i_0 i_1 \cdots i_j} e_{i_0} \otimes e_{i_{\sigma(1)}} \otimes \cdots \otimes e_{i_{\sigma(j)}}$. The map σ induces a diffeomorphism $P(\boldsymbol{R}^l \otimes \otimes^j_{i=1} \boldsymbol{R}^{k_i}) \to P(\boldsymbol{R}^l \otimes \otimes^j_{i=1} \boldsymbol{R}^{\sigma(k_i)})$. The group $PL(l) \times \prod^j_{i=1} PL(k_{\sigma(i)})$ naturally acts on $P(\boldsymbol{R}^l \otimes \otimes^j_{i=1} \boldsymbol{R}^{\sigma(k_i)})$, and the action satisfies the condition $(g, A_{\sigma(1)}, \dots, A_{\sigma(j)}) \cdot \sigma(w) = \sigma((g, A_1, \dots, A_j) \cdot w)$. It is easy to prove the following.

Proposition 6.1. *Assume that a point* w *in* $P(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i})$ *gives an open orbit of* $PL(l) \times \prod_{i=1}^j PL(k_i)$. *Then we have the following:*

(1) for any permutation σ *of* $\{1, 2, \cdots j\}$ *the product group* $PL(l) \times \prod_{i=1}^j PL(k_{\sigma(i)})$ *admits an open orbit given by* $\sigma(w)$. *Isotropy subgroups* $PL(l) \times \prod_{i=1}^j PL(k_i)_w$ *and* $PL(l) \times \prod_{i=1}^j PL(k_{\sigma(i)})_{\sigma(w)}$ *are isomorphic.*

(2) The product group $PL(l) \times \prod_{i=1}^j PL(k_i) \times PL(1)$ *acts on* $P(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i} \otimes \boldsymbol{R})$, *which is identified with* $P(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i})$ *naturally. Via this identification* w *gives an open orbit of* $PL(l) \times \prod_{i=1}^j PL(k_i) \times PL(1)$, *and isotropy subgroups* $PL(l) \times \prod_{i=1}^j PL(k_i)_w$ *and* $PL(l) \times \prod_{i=1}^j PL(k_i) \times PL(1)_w$ *are isomorphic.*

From the Lie group $PL(l) \times \prod_{i=1}^j PL(k_i)$ we can obtain several new Lie groups by Propositions 6.1 and castling transformations. For example let L be a Lie subgroup of $PL(3)$ such that L admits an open orbit $L.x$ in $P(\boldsymbol{R}^3)$. Then we obtain the sequence of new groups $*L \times PL(2)$, $L \times PL(2) \times PL(5)$, $*L \times PL(5) \times PL(13)$, $L \times PL(5) \times PL(13) \times PL(194)$, which admit open orbits in projective spaces. Note that $*L \times PL(2)$ is regarded as a subgroup $*L \otimes PL(2)$ of $PL(6)$. Next we apply Proposition 6.1 to Cartan connections.

Proposition 6.2. *Let* Q *be a manifold equipped with a* $\mathfrak{g}$*-valued 1-from* ω. *Assume that* $(Q \times \prod_{i=1}^j PL(k_i), \omega \times \prod_{i=1}^j \Lambda_1)$ *gives a Cartan connection over a manifold* N *of type*

$$PL(l) \times \prod_{i=1}^j PL(k_i) / PL(l) \times \prod_{i=1}^j PL(k_i)_w,$$

which is a subgeometry of $PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i}) / PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i})_w$ *given by* $\imath : PL(l) \times \prod_{i=1}^j PL(k_i) \hookrightarrow PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i})$. *Then*

1. *for any permutation* σ *of* $\{1, 2, \cdots j\}$, *the pair* $(Q \times \prod_{i=1}^j PL(k_{\sigma(i)}), \omega \times \prod_{i=1}^j \Lambda_1)$ *gives a Cartan connection over* N *of type*

$$PL(l) \times \prod_{i=1}^j PL(k_{\sigma(i)}) / PL(l) \times \prod_{i=1}^j PL(k_{\sigma(i)})_{\sigma(w)},$$

   which is a subgeometry of $PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_{\sigma(i)}}) / PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_{\sigma(i)}})_{\sigma(w)}$.

2. $(Q \times \prod_{i=1}^j PL(k_i) \times PL(1), \omega \times \prod_{i=1}^j \Lambda_1 \times \Lambda_1)$ *gives a Cartan connection over a manifold* N *of type*

$$PL(l) \times \prod_{i=1}^j PL(k_i) \times PL(1) / PL(l) \times \prod_{i=1}^j PL(k_i) \times PL(1)_w,$$

   which is a subgeometry of $PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i} \otimes \boldsymbol{R}) / PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i} \otimes \boldsymbol{R})_w$.

We consider the castling transformation of $(Q \times \prod_{i=1}^j PL(k_i), \omega \times \prod_{i=1}^j \Lambda_1)$. The group $PL(l) \times \prod_{i=1}^j PL(k_i)$ can be identified with a subgroup of $PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i})$ by the map

$F : (g \times A_1 \times \cdots \times A_j) \mapsto g \otimes A_1 \otimes \cdots \otimes A_j$. Then one form $\omega \times \prod_{i=1}^{j} \Lambda_1$ can be identified with the one form $dF \circ \omega \times \prod_{i=1}^{j} \Lambda_1$, which is computed as follows:

$$\begin{aligned} dF \circ \omega \times \prod_{i=1}^{j} \Lambda_1(X, Y_1, \cdots, Y_j) \quad &= \quad \omega(X) \otimes I \otimes \cdots \otimes I + I \otimes Y_1 \otimes \cdots \otimes Y_j \\ & \qquad + I \otimes \cdots \otimes I \otimes Y_j. \end{aligned}$$

Thus the group isomorphism $*$ of $PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^{j} \boldsymbol{R}^{k_i})$ is restricted to the subgroup $PL(l) \times \prod_{i=1}^{j} PL(k_i)$ by

$$* : (g, A_1, \ldots, A_j) \mapsto (*g, *A_1, \ldots, *A_j).$$

The Lie algebra isomorphism $*$ of $\mathfrak{sl}(\boldsymbol{R}^l \otimes \otimes_{i=1}^{j} \boldsymbol{R}^{k_i})$ is restricted to $\mathfrak{g} \times \prod_{i=1}^{j} \mathfrak{sl}(k_i)$ and we have

$$*(\omega \times \prod_{i=1}^{j} \Lambda_1) = *\omega \times \prod_{i=1}^{j} *\Lambda_1.$$

By using Propositions 3.3 and 6.2, we can apply castling transformations successively. For example let us consider a projective Cartan connection (Q, ω) over 2-dimensional manifold M. The bundle Q is a principal fiber bundle over M with structure group $PL(3)_v$, where v is an element of $P(\boldsymbol{R}^3)$ and ω is a $\mathfrak{sl}(3)$-valued 1-form. Then for example we obtain the following sequence by successive castling transformations:

$$\begin{aligned} &(1)\ (Q, \omega) \longrightarrow (2)\ (Q, *\omega) \longrightarrow (3)\ (Q \times PL(2), *\omega \times \Lambda_1) \\ &\longrightarrow (4)\ (Q \times PL(2), \omega \times *\Lambda_1) \longrightarrow (5)\ (Q \times PL(2) \times PL(5), \omega \times *\Lambda_1 \times \Lambda_1). \end{aligned}$$

Now consider the Grassmannian manifold $PL(l)/PL(l)_{<v>}$, and we denote by ρ the isotropy representation of $PL(l)_{<v>}$. Then ρ takes values in $GL(n) \otimes GL(m)$. When $\rho(g)$ is expressed as $A \otimes B$, we define a projective linear representation $\rho_m : PL(l)_{<v>} \to PL(m)$ by $\rho_m(g) = \overline{B}$. The isotropy group $PL(l) \times PL(m)_v$ is equal to the set $\{(g, \rho_m(g)) \mid g \in PL(l)_{<v>}\}$. Thus the two isotropy groups $PL(l)_{<v>}$ and $PL(l) \times PL(m)_v$ are isomorphic.

Here we omitted the process of 1 and 2 in Proposition 6.2. In detail we omitted the Cartan connections $(Q \times PL(1), \omega \times \Lambda_1)$ and $(Q \times PL(2) \times PL(1), *\omega \times \Lambda_1 \times \Lambda_1)$. In the above process we choose a point $v^\perp \in V_{2,3}$ and identify $v^\perp$ with a point w of $P(\boldsymbol{R}^3 \otimes \boldsymbol{R}^2)$, and choose a point $w^\perp \in V_{5,3\cdot 2}$. Then the structure group of each Cartan connection is given by

$$\begin{aligned} &(1) \quad PL(3)_v, \\ &(2) \quad PL(3)_{<v^\perp>} = *PL(3)_v, \\ &(3) \quad PL(3) \times PL(2)_{v^\perp} = \{(*g, \rho_2(*g)) \mid g \in PL(3)_v\}, \\ &(4) \quad PL(3) \times PL(2)_{<w^\perp>} = \{(g, *\rho_2(*g))\}, \\ &(5) \quad PL(3) \times PL(2) \times PL(5)_{w^\perp} = \{(g, *\rho_2(*g), \rho_5(g \otimes *\rho_2(*g)))\}. \end{aligned}$$

We express the action of $PL(2)_v$ on Q as $u \cdot g = ug$ for $u \in Q$ and $g \in PL(2)_v$. Then each structure group acts on the bundle of each Cartan connection as follows:

$$\begin{aligned} &(2) \quad u \cdot *g = ug, \\ &(3) \quad (u, A) \cdot (*g, \rho_2(*g)) = (ug, A\rho_2(*g)), \\ &(4) \quad (u, A) \cdot (g, *\rho_2(*g)) = (ug, A\rho_2(*g)), \\ &(5) \quad (u, A, B) \cdot (g, *\rho_2(*g), \rho_5(g \otimes *\rho_2(*g))) = (ug, A\rho_2(*g), B\rho_5(g \otimes *\rho_2(*g))). \end{aligned}$$

Successive castling transformations yields a sequence of manifolds admitting a projective structure or a Grassmannian structure. From now on we characterize those manifolds.

Lemma 6.3. *Let v be a point in $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^\alpha)$ and assume that $PL(l) \times PL(\alpha) \cdot v$ gives an open orbit in $P(\boldsymbol{R}^l \otimes \boldsymbol{R}^\alpha)$. Suppose that a Lie group G is obtained by successive castling transformations from $PL(l) \times PL(\alpha)$, and a point w in $P(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i})$ is obtained from v. Then G can be written as $PL(l) \times \prod_{i=1}^j PL(k_i)$ and $G \cdot w$ gives an open orbit in $P(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i})$ again. Moreover there exists a projective linear representation $f_i\colon PL(l)_{<v>} \to PL(k_i)$ $(0 \leq i \leq j)$ such that $G_w = \{(f_0(g), f_1(g), \ldots, f_j(g)) \mid g \in PL(l)_{<v>}\}$, where $f_0 = id$ or $*$. Thus G_w is isomorphic to $PL(l)_{<v>}$.*

Proof. First about the group $PL(l) \times PL(\alpha)$, the element of the isotropy subgroup $PL(l) \times PL(\alpha)_v$ is expressed as $(g, \rho_\alpha(g))$, where ρ_α is the projective linear representation of $PL(l)$ introduced after Proposition 3.1. We now proceed by induction. Assume that there exists $w \in P(\boldsymbol{R}^l \otimes \otimes_{i=1}^j \boldsymbol{R}^{k_i})$ such that the group $PL(l) \times \prod_{i=1}^j PL(k_i)$ admits an open orbit given by w, and an element of the isotropy group $PL(l) \times \prod_{i=1}^j PL(k_i)_w$ is expressed as $(f_0(g), f_1(g), \ldots, f_j(g))$ for some $g \in PL(l)_{<v>}$. Then the point w must belong to $V_{k_j, lk_1 \cdots k_{j-1}}$. A group obtained by using the assertion (1) or (2) in Proposition 6.1 also admits an open orbit and its isotropy group is described by using the projective linear representations of $PL(l)_{<v>}$. Since successive castling transformation consists of Propositions 6.1 and 3.1, it is enough to consider the effect of castling transformation described in Proposition 3.1. By castling transformation of the group $PL(l) \times \prod_{i=1}^j PL(k_i)$, we obtain the group $PL(l) \times \prod_{i=1}^{j-1} PL(k_i) \times PL(lk_1 \cdots k_{j-1} - k_j)$ and a fixed pt $w^\perp \in V_{lk_1 \cdots k_{j-1} - k_j, lk_1 \cdots k_{j-1}}$, which gives an open orbit in $P(\boldsymbol{R}^l \otimes \otimes_{i=1}^{j-1} \boldsymbol{R}^{k_i} \otimes \boldsymbol{R}^{lk_1 \cdots k_{j-1} - k_j})$. Here we regard $PL(l) \times \prod_{i=1}^{j-1} PL(k_i)$ as the subgroup of $PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^{j-1} \boldsymbol{R}^{k_i})$. Then about the isotropy group $PL(l) \times \prod_{i=1}^{j-1} PL(k_i) \times PL(lk_1 \cdots k_{j-1} - k_j)_{w^\perp}$ its element is expressed as $(*f_0(g), *f_1(g), \ldots, *f_{j-1}(g), \rho_{lk_1 \cdots k_{j-1} - k_j}(*f_0(g) \otimes *f_1(g) \otimes \cdots \otimes *f_{j-1}(g)))$ for some $g \in PL(l)_{<v>}$, where $\rho_{lk_1 \cdots k_{j-1} - k_j}$ is the projective linear representation of $PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^{j-1} \boldsymbol{R}^{k_i})_{<w^\perp>}$. Thus we completes the induction step. ☐

Now let (Q, ω) be a Grassmannian Cartan connection of type $(l - \alpha, \alpha)$ over M. Thus the model space is $PL(l)/PL(l)_{<v>}$, where $< v >$ is an element of $Gr_{\alpha,l}$. By successive castling transformations from (Q, ω) we obtain a Cartan connection $(Q \times \prod_{i=1}^j PL(k_i), \omega' \times \prod_{i=1}^j (\Lambda_1)_i)$ of type $PL(l) \times \prod_{i=1}^j PL(k_i)/\ PL(l) \times \prod_{i=1}^j PL(k_i)_w$, where $(\Lambda_1)_i = \Lambda_1$ or $*\Lambda_1$ and $\omega' = \omega$ or $\omega' = *\omega$. We denote by N the base space of $(Q \times \prod_{i=1}^j PL(k_i), \omega' \times \prod_{i=1}^j (\Lambda_1)_i)$. Then the Cartan connection $(Q \times \prod_{i=1}^j PL(k_i), \omega' \times \prod_{i=1}^j (\Lambda_1)_i)$ over N induces a projective Cartan connection over N. The next proposition determines the relation of base spaces obtained by successive castling transformations. Assume that $k_i \neq 1$. Remove the s-th component from $(Q \times \prod_{i=1}^j PL(k_i), \omega' \times \prod_{i=1}^j (\Lambda_1)_i)$ and denote it by $(Q \times \prod_{i \neq s}^j PL(k_i), \omega' \times \prod_{i \neq s}^j (\Lambda_1)_i)$.

Proposition 6.4. *Choose s and t satisfying* $1 \leq s < t \leq j$.
(1) *The base space N is a principal fiber bundle over M with group $\prod_{i=1}^j PL(k_i)$.*
(2) *The pair $(Q \times \prod_{i \neq s}^j PL(k_i), \omega' \times \prod_{i \neq s}^j (\Lambda_1)_i)$* (resp. $(Q \times \prod_{i \neq t}^j PL(k_i), \omega' \times \prod_{i \neq t}^j (\Lambda_1)_i)$) is a Cartan connection over a manifold L (resp. K), which is a subgeometry of a Grassmannian Cartan connection of type $(lk_1 \cdots k_{s-1} k_{s+1} \cdots k_j - k_s, k_s)$ (resp. $(lk_1 \cdots k_{t-1} k_{t+1} \cdots k_j - k_t, k_t)$). The base space L is a $\prod_{i \neq s}^j PL(k_i)$-bundle over M.

(3) The base space N is regarded as a principal fiber bundle over L (resp. K) with group $PL(k_s)$ (resp. $PL(k_t)$), and $PL(k_t)$ acts on N and L. Thus we obtain following diagram:

$$\begin{tikzcd}
 & PL(k_s) \arrow[r] & N \arrow[dll] \arrow[drr] & PL(k_t) \arrow[l] & \\
L & PL(k_t) \arrow[l] & & PL(k_s) \arrow[r] & K.
\end{tikzcd}$$

Moreover N is isomorphic to the bundle $P_tL/GL(lk_1 \cdots k_{s-1}k_{s+1} \cdots k_j - k_s) \otimes GL(1)$ and the action of $PL(k_t)$ on L induces the action on N by the differential. The quotient $N/PL(k_t)$ is isomorphic to K.

Proof. We can assume that the number t is equal to j without loss of generality. Firstly we show that N is a principal fiber bundle over M. From the assumption N is diffeomorphic to the quotient manifold $Q \times \prod_{i=1}^{j} PL(k_i)/PL(l) \times \prod_{i=1}^{j} PL(k_i)_w$, thus we identify N with this quotient manifold. By using this identification the group $\prod_{i=1}^{j} PL(k_i)$ naturally acts on N as follows: let $[z, A_1, \ldots, A_j]$ be an element of N and $(B_1, \ldots, B_j)$ be an element of $\prod_{i=1}^{j} PL(k_i)$. Then we define the action by $[z, A_1, \ldots, A_j] \cdot (B_1, \ldots, B_j) := [z, B_1^{-1}A_1, \ldots, B_j^{-1}A_j]$. This action is free. By using the projection $\pi : Q \to M$ we define the projection $\pi_N : N \to M$ by $[z, A_1, \ldots, A_j] \mapsto \pi(z)$. Moreover $\pi : Q \to M$ is a principal fiber bundle, thus for each open neighborhood U of M there is a local trivialization $\pi^{-1}(U) \to U \times PL(l)_{<v>}$ mapping z to $(\pi(z), \phi(z))$. Now by Lemma 6.3 the isotropy subgroup $PL(l) \times \prod_{i=1}^{j} PL(k_i)_w$ is given by the set $\{(f_0(g), f_1(g), \ldots, f_j(g)) \mid g \in PL(l)_{<v>}\}$. We define the map $\Phi : \pi_N^{-1}(U) \to U \times \prod_{i=1}^{j} PL(k_i)$ by

$$[z, A_1, \ldots, A_j] \mapsto (\pi(z), {f_1}'(\phi(z))A_1^{-1}, \ldots, {f_j}'(\phi(z))A_j^{-1}),$$

where ${f_i}' = f_i$ when $(\Lambda_1)_i = \Lambda_1$ or ${f_i}' = *f_i$ when $(\Lambda_1)_i = *\Lambda_1$. The map Φ is well-defined and diffeomorphism, moreover preserving the action of $\prod_{i=1}^{j} PL(k_i)$. By using π_N and the local trivializations Φ, it follows that N has a structure of principal fiber bundle over M with group $\prod_{i=1}^{j} PL(k_i)$.

Since $\prod_{i=1}^{j-1}\{e\} \times PL(k_j)$ is a normal closed subgroup of structure group $\prod_{i=1}^{j} PL(k_i)$ of N, the quotient $N/PL(k_j)$ is again a principal fiber bundle over M with group $\prod_{i=1}^{j-1} PL(k_i)$. Put $H = \prod_{i=1}^{j-1} PL(k_i)$. Then the manifold $Q \times H$ is regarded as a principal fiber bundle over M with group $PL(l)'_{<v>} \times H$, where $PL(l)'_{<v>} = PL(l)_{<v>}$ if $\omega' = \omega$ or $PL(l)'_{<v>} = *PL(l)_{<v>}$ if $\omega' = *\omega$. The group $PL(l)'_{<v>} \times H$ contains the closed subgroup $PL(l) \times H_{<w>}$. Hence we obtain the fiber bundle $Q \times H$ over the quotient manifold $Q \times H/PL(l) \times H_{<w>}$ with structure group $PL(l) \times H_{<w>}$. There is a diffeomorphism from $Q \times H/PL(l) \times H_{<w>}$ to $N/PL(k_j)$ defined by $[u, A_1, \ldots, A_{j-1}] \mapsto [u, A_1, \ldots, A_{j-1}, e]PL(k_j)$. By Proposition 3.3 $(Q \times H, \omega \times \prod_{i=1}^{j-1} \Lambda_1)$ is a Cartan connection over $N/PL(k_j)$ of type $PL(l) \times H/PL(l) \times H_{<w>}$. From the assumption the model space $PL(l) \times H/PL(l) \times H_{<w>}$ is a subgeometry of the Grassmannian manifold $PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^{j-1}\boldsymbol{R}^i)/PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^{j-1}\boldsymbol{R}^i)_{<w>}$ according to the map $F : PL(l) \times H \hookrightarrow PL(\boldsymbol{R}^l \otimes \otimes_{i=1}^{j-1}\boldsymbol{R}^i)$ defined by the tensor product. Note that a point w is included in $V_{k_j, lk_1 \cdots k_{j-1}}$. Therefore $N/PL(k_j)$ admits a Grassmannian Cartan connection of type $(lk_1 \cdots k_{j-1} - k_j, k_j)$.

We observe naturally $PL(k_s)$ acts freely on $Q \times H$ by $(u, A_1, \ldots, A_s, \ldots, A_{j-1}) \cdot B_s := (u, A_1, \ldots, B_s^{-1}A_s, \ldots, A_{j-1})$ for $B_s \in PL(k_s)$. We denote this right action by R_{B_s}. Put

$L = N/PL(k_j)$. The action of $PL(k_s)$ on $Q \times \prod_{i=1}^{j-1} PL(k_i)$ induces the action of $PL(k_s)$ on L by using the bundle isomorphism between L and $Q \times H/PL(l) \times H_{<w>}$.

We fix the complementary subspace $\mathfrak{m}$ of $\mathfrak{sl}(l)_{<v>}$ in $\mathfrak{sl}(l)$, thus we have $\mathfrak{sl}(l) = \mathfrak{m} \oplus \mathfrak{sl}(l)_{<v>}$. Then $\mathfrak{m}' \times \prod_{i=1}^{j-1} \mathfrak{sl}(k_i)$ gives a complementary subspace of $\mathfrak{sl}(l) \times \prod_{i=1}^{j-1} \mathfrak{sl}(k_i)_{<w>}$ in $\mathfrak{sl}(l) \times \prod_{i=1}^{j-1} \mathfrak{sl}(k_i)$, where $\mathfrak{m}' = \mathfrak{m}$ when $\omega' = \omega$ or $\mathfrak{m}' = *\mathfrak{m}$ when $\omega' = *\omega$. Furthermore the differential dF of F induces a linear isomorphism $\widehat{dF}$ from $\mathfrak{m}' \times \prod_{i=1}^{j-1} \mathfrak{sl}(k_i)$ to $M(lk_1 \cdots k_{j-1} - k_j, k_j)$.

We denote by ρ the isotropy representation of the model space $PL(l) \times H/PL(l) \times H_{<w>}$. Denote $Q \times H$ by P, and the natural projection $P \to P/ker\rho$ by ρ. We denote by $\widetilde{P}$ the quotient space $P/ker\rho$. The action of $PL(k_s)$ on P induces the action on $\widetilde{P}$. Then there exists a unique 1-form θ on $\widetilde{P}$ such that $\rho^*\theta = \omega' \times \prod_{i=1}^{j-1} \Lambda_1{}_{\mathfrak{m}' \times \prod_{i=1}^{j-1} \mathfrak{sl}(k_i)}$. The pair $(\widetilde{P}, \theta)$ gives a $\rho(PL(l) \times H_{<w>})$-structure over L, which is a subbundle of a $GL(lk_1 \cdots k_{j-1} - k_j) \otimes GL(k_j)$-structure P_tL. By Proposition 2.5 the imbedding ι of $\widetilde{P}$ into P_tL is given by the restriction of the bundle isomorphism $t : \mathcal{L}(L) \to \mathcal{L}(L)$, which is defined by $t : x \mapsto x \circ \widehat{dF}^{-1}$.

The equality ${R_{B_s}}^*\omega \times \prod_{i=1}^{j-1} \Lambda_1 = \omega \times \prod_{i=1}^{j-1} \Lambda_1$ yields ${R_{B_s}}^*\rho^*\theta = \rho^*\theta$. Since we have the commutative diagram

$$\begin{tikzcd}
P \arrow[r, "R_{B_s}"] \arrow[d, "\rho"'] \arrow[dr, phantom, "\circlearrowleft"] & P \arrow[d, "\rho"'] \\
\widetilde{P} \arrow[r, "R_{B_s}"] \arrow[d] \arrow[dr, phantom, "\circlearrowleft"] & \widetilde{P} \arrow[d] \\
L \arrow[r, "R_{B_s}"] & L,
\end{tikzcd} \tag{6.1}$$

it follows that ${R_{B_s}}^*\rho^*\theta = \rho^*{R_{B_s}}^*\theta$. Thus ${R_{B_s}}^*\theta = \theta$. Therefore the action $R_{B_s} : \widetilde{P} \to \widetilde{P}$ is induced by the differential of the action $R_{B_s} : L \to L$. Moreover $R_{B_s} : \tilde{P} \to \tilde{P}$ is uniquely extended to the action $R_{B_s} : P_tL \to P_tL$ by

$$\iota \circ \rho(z, A_1, \ldots, A_s, \ldots, A_{j-1})B \otimes C \cdot D = \iota \circ \rho(z, A_1, \ldots, D^{-1}A_s, \ldots, A_{j-1})B \otimes C,$$

where $(z, A_1, \ldots, A_s, \ldots, A_{j-1}) \in Q'$, $B \otimes C \in GL(lk_1 \cdots k_{j-1} - k_j) \otimes GL(k_j)$ and $D \in PL(k_s)$. This action naturally induces the action of $PL(k_s)$ on $P_tL/GL(lk_1 \cdots k_{j-1} - k_j) \otimes GL(1)$. We describe the process that the action of $PL(k_s)$ on P induces the actions on other manifolds by the following diagram.

$$\begin{tikzcd}
PL(k_s) \arrow[d, bend left] & PL(k_s) \arrow[d, bend left] & PL(k_s) \arrow[d, bend left] & PL(k_s) \arrow[d, bend left] \\
P \arrow[r, dashed] & \widetilde{P} \arrow[r, dashed] & P_tL \arrow[r, dashed] & P_tL/GL(lk_1 \cdots k_{j-1} - k_j) \otimes GL(1)
\end{tikzcd}$$

By the proof of Proposition 3.3 there is a $PL(k_j)$-bundle isomorphism from $N = P \times PL(k_j)/H \times \prod_{i=1}^{j} PL(k_i)_w$ to $P_tL/GL(lk_1 \cdots k_{j-1} - k_j) \otimes GL(1)$. Via this isomorphism we see that the action of $PL(k_s)$ on N coincides with the one on $P_tL/GL(lk_1 \cdots k_{j-1} - k_j) \otimes GL(1)$. Thus the induced action of $PL(k_s)$ on N is given by $[u, A_1, \ldots, A_s, \ldots, A_j] \cdot B_s = [u, A_1, \ldots, B_s^{-1}A_s, \ldots, A_j]$ for $B_s \in PL(k_s)$. Consequently the action of $PL(k_s)$ on L induces the action on N by the differential. By this action we have the quotient $K = N/PL(k_s)$. On the other hand let σ be a permutation defined by

$$\sigma = \begin{pmatrix} 1 & 2 & \cdots & s-1 & s & s+1 & \cdots & j-1 & j \\ 1 & 2 & \cdots & s-1 & s+1 & \cdots & j-1 & j & s \end{pmatrix}.$$

Then by Proposition 6.2 the given Cartan connection $(Q \times \prod_{i=1}^{j} PL(k_i), \omega' \times \prod_{i=1}^{j} (\Lambda_1)_i)$ over N can be regarded as a Cartan connection over N of type $PL(l) \times \prod_{i=1}^{s-1} PL(k_i) \times \prod_{i=s+1}^{j} PL(k_i) \times PL(k_s)/PL(l) \times \prod_{i=1}^{s-1} PL(k_i) \times \prod_{i=s+1}^{j} PL(k_i) \times PL(k_s)_{\sigma(w)}$, which is a subgeometry of the projective space $P(\boldsymbol{R}^l \otimes \otimes_{i=1}^{s-1} \boldsymbol{R}^{k_i} \otimes \otimes_{i=s+1}^{j} \boldsymbol{R}^{k_i} \otimes \boldsymbol{R}^s)$. By Proposition 3.3 it follows that K admits a Grassmannian Cartan connection of type $(lk_1 \cdots k_{s-1}k_{s+1} \cdots k_j - k_s, k_s)$. □

7 Examples of successive castling transformations

7.1

Let $(P_L, [\chi])$ be a projective structure over n-dimensional manifold M, and (Q, ω) be a projective Cartan connection over M constructed from $(P_L, [\chi])$. As we demonstrate it after Proposition 6.2 by successive castling transformations we can obtain the following Cartan connections:

$$\begin{aligned} &(Q, \omega) \longrightarrow (Q \times PL(n), *\omega \times \Lambda_1) \longrightarrow (Q \times PL(n) \times PL(n^2+n-1), \omega \times *\Lambda_1 \times \Lambda_1) \\ &\longrightarrow (Q \times PL(n^2+n-1), \omega \times \Lambda_1). \end{aligned}$$

We denote the base space of (Q, ω) by M_1, the one of $(Q \times PL(n), *\omega \times \Lambda_1)$ by M_n and so on. Generally $M_{k_1 \times \cdots \times k_j}$ denotes a principal fiber bundle over M with structure group $\prod_{i=1}^{j} PL(k_i)$. Then following the above successive Cartan connections from (Q, ω) we obtain the sequence of base spaces: $M_1 \longrightarrow M_2 \longrightarrow M_{2\times 5} \longrightarrow M_5$. The Cartan connection $(Q \times PL(n), *\omega \times \Lambda_1)$ over M_n and $(Q \times PL(n) \times PL(n^2+n-1), \omega \times *\Lambda_1 \times \Lambda_1)$ over $M_{n\times(n^2+n-1)}$ induce projective Cartan connections, and the Cartan connection $(Q \times PL(n^2+n-1), \omega \times \Lambda_1)$ over M_{n^2+n-1} induces a Grassmannian Cartan connection. Now we describe those base spaces more explicitly:

Proposition 7.1. *M_n is isomorphic to the projective frame bundle of M, and $M_{n\times(n^2+n-1)}$ and M_{n^2+n-1} is isomorphic to the following bundles $\widetilde{\mathcal{L}}_{n\times(n^2+n-1)}$ and $\widetilde{\mathcal{L}}_{n^2+n-1}$:*

$$\{(u_1, u_2) \mid u_1 : \textit{projective frame of } T_pM, u_2 : \textit{projective frame of } T_pM \rtimes \mathfrak{sl}(T_pM), p \in M\},$$

$$\{(u_2) \mid u_2 : \textit{projective frame of } T_pM \rtimes \mathfrak{sl}(T_pM), p \in M\}.$$

Proof. By definition Q is a principal fiber bundle over M with structure group $PL(n)_v$, where $v = (1, 0, \cdots, 0)$ is an element of $V_{1,n}$. From the construction we have the injection $h : \mathcal{L}(M) \hookrightarrow Q$ corresponding to the injection $\iota : GL(n) \hookrightarrow PL(n+1)_v$, which is defined by

$$\iota : A \mapsto \begin{pmatrix} 1 & 0 \\ 0 & A \end{pmatrix}.$$

By castling transformation of (Q, ω) we obtain the Cartan connection $(Q \times PL(n), *\omega \times \Lambda_1)$ over M_n whose structure group is $PL(n+1) \times PL(n)_{<v^\perp>}$, where $v^\perp = \begin{pmatrix} 0 \\ I_n \end{pmatrix}$ is an element of $V_{n,n+1}$.

Then $\iota(GL(n))\times PL(n)_{v^\perp}$ is given by the set $\{(*\iota(A),\bar{A})\}$. The manifold $h(\mathcal{L}(M))\times PL(n)$ is a principal fiber bundle over M_n with structure group $\iota(GL(n))\times PL(n)_{v^\perp}$, and gives a reduction of $Q\times PL(n)$. Then we have the following bundle isomorphism $h(\mathcal{L}(M))\times PL(n)/\iota(GL(n))\times PL(n)_{v^\perp}\to\mathcal{L}(M)/GL(1)$ defined by

$$[h(x),g]\mapsto q(x)g^{-1},$$

where q is the projection $\mathcal{L}(M)\to\mathcal{L}(M)/GL(1)$. Note that the action of $\iota(GL(n))\times PL(n)_{v^\perp}$ on $h(\mathcal{L}(M))\times PL(n)$ is given by $(h(x),g)\cdot(*\iota(A),\bar{A})=(h(x)\iota(A),g\bar{A})$. Thus the base space M_n is isomorphic to the projective frame bundle of M. Concerning the Cartan connection $(Q\times PL(n)\times PL(n^2+n-1),\omega\times *\Lambda_1\times\Lambda_1)$ over $M_{n\times(n^2+n-1)}$, the structure group of $Q\times PL(n)\times PL(n^2+n-1)$ is $PL(n+1)\times PL(n)\times PL(n^2+n-1)_{w^\perp}$, where $w^\perp$ is a fixed projective frame of the vector space $\left(\begin{array}{c}M(1,n)\\ \hline \mathfrak{sl}(n)\end{array}\right)$. When we are given a base x of a vector space V, we denote by $q(x)$ the projective frame, then q gives the projection from the linear Stiefel manifold to the projective Stiefel manifold corresponding to the projection $-:GL(n)\to PL(n)$. Then the subgroup $\iota(GL(n))\times PL(n)\times PL(n^2+n-1)_{w^\perp}$ is given by the set $\{(\iota(A),*\bar{A},\rho_{n^2+n-1}(\iota(A)\otimes *\bar{A}))\}$, where

$$\rho_{n^2+n-1}(\iota(A)\otimes *\bar{A})=\overline{\begin{pmatrix}A & 0\\ 0 & Ad(A)\end{pmatrix}}$$

with respect to the basis

$$\{\left(\begin{array}{c}e_i\\ \hline 0\end{array}\right)(1\leq i\leq n),\left(\begin{array}{c}0\\ \hline E^k_j-\delta^k_jE^n_n\end{array}\right)(1\leq i,j\leq n)\}.$$

With respect to a base $x=\{X_1,\cdots,X_n\}$ of T_pM and $Y=\sum Y^i_jE^j_i\in\mathfrak{sl}(n)$ we define an element $Y_x\in\mathfrak{sl}(T_pM)$ by $Y_x:=x\circ Y\circ x^{-1}$, where we regard x as a linear isomorphism. That is to say Y_x is the map $Y_x:X_k\mapsto\sum Y^i_kX_i$. Then for $A\in GL(n)$ we have $Y_{xA}=Ad(A)(Y_x)$. We denote by ${E'}^k_j$ the element $E^k_j-\delta^k_jE^n_n$ of $\mathfrak{sl}(n)$. Now we define the map $\phi:h(\mathcal{L}(M))\times PL(n)\times PL(n^2+n-1)\to\widetilde{\mathcal{L}}_{n\times(n^2+n-1)}$ by

$$[h(x),g_1,g_2]\mapsto\{q(x){g_1}^{-1},q(x,({E'}^k_j)_x){g_2}^{-1})\}$$

We show that the map ϕ is well defined. We put

$$(h(x'),g'_1,g'_2):=(h(x),g_1,g_2)\cdot(\iota(A),*\bar{A},\rho_{n^2+n-1}(\iota(A)\otimes *\bar{A})),$$

which is equal to $(h(xA),g_1\bar{A},g_2\rho_{n^2+n-1}(\iota(A)\otimes *\bar{A}))$. Then we have

$$\begin{aligned}&\{q(x'){g'_1}^{-1},q(x',({E'}^k_j)_{x'}){g'_2}^{-1})\}\\ =\ &\{q(xA)(g_1\bar{A})^{-1},q(xA,({E'}^k_j)_{xA})(g_2\rho_{n^2+n-1}(\iota(A)\otimes *\bar{A}))^{-1}\}.\end{aligned}$$

The last expression is equal to $\{q(x){g_1}^{-1},q(x,({E'}^k_j)_x){g_2}^{-1})\}$ since we have

$$q(xA,({E'}^k_j)_{xA})=q(x,({E'}^k_j)_x)\rho_{n^2+n-1}(\iota(A)\otimes *\bar{A}).$$

Moreover ϕ is a bundle isomorphism. Likewise we can show that M_{n^2+n-1} is isomorphic to $\widetilde{\mathcal{L}}_{n^2+n-1}$. $\square$

Now we explain how these manifolds $M_{k_1\times\cdots\times k_j}$ are related in the case of $n=2$. From Propositions 7.1 and 3.3 M_2 is a projective frame bundle of M_1 and $M_{2\times 5}$ is a projective frame bundle of M_2. Thus M_2 has the right action of $PL(2)$ and this action gives rise to the action of $PL(2)$ on the frame bundle of M_2 by the differential. Moreover $PL(2)$ naturally acts on the projective frame bundle of M_2, which is equal to $M_{2\times 5}$. With respect to this action of PL(2) the quotient $M_{2\times 5}/PL(2)$ is equal to M_5 from Proposition 6.4. Furthermore (Q,ω), $(Q\times PL(2), *\omega\times\Lambda_1)$ and $(Q\times PL(2)\times PL(5), \omega\times *\Lambda_1\times\Lambda_1)$ induces projective Cartan connections and $(Q\times PL(5), \omega\times\Lambda_1)$ induces Grassmannian Cartan connections. The same result holds true about the general dimension n. If we continue the successive castling transformations we can obtain the following tree, where we only describe the base spaces. We abbreviate $M_{k_1\times\cdots\times k_j}$ to $k_1\times\cdots\times k_j$. If a Cartan connection over a base space induces a Grassmannian structure of type (β,α) then we write $GL(\beta)\otimes GL(\alpha)$ under the base space. If a Cartan connection induces a projective structure, then we write nothing.



The relation of the base spaces of the tree is completely described by Proposition 6.4. Moreover for any manifold M the all principal fiber bundles over M underlined in blue admit a Grassmannian structure. If the given projective structure on M is projectively flat, then the all principal fiber bundle over M underlined in red admit a flat projective structure and manifolds underlined in blue admit a flat Grassmannian Cartan connection. Especially a manifold $M_{k_1\times\cdots\times k_{j-1}}$ underlined in blue admits a Grassmannian structure of type (β, α) which is given by an extension of the bundle $\mathcal{L}(M) \times \prod_{i=1}^{j-1} PL(k_i)$ over $M_{k_1\times\cdots\times k_{j-1}}$ whose structure group is isomorphic to $GL(2)$. This $GL(2)$-bundle gives a reduction of $GL(3k_1 \cdots k_{j-1} - k_j) \otimes GL(k_j)$-structure on $M_{k_1\times\cdots\times k_{j-1}}$. We can prove this assertion generally as follows: we use the same notations in Proposition 6.4, thus M admits a Grassmannian Cartan connection of type $(l-\alpha, \alpha)$, and N is a fiber bundle $M_{k_1\times\cdots\times k_j}$ equipped with a projective structure. Denote by L the quotient $N/PL(k_j)$ and by P the bundle $Q \times H$ over L with structure group $PL(l) \times H_{<w>}$. Now we assume that $lk_1 \cdots k_{j-1} - k_j \neq 1$, hence L admits a Grassmannian structure P_tL. The Lie group homomorphism $\rho : PL(l) \times H_{<w>} \to GL(M(l-\alpha, \alpha) \times \prod_{i=1}^{j-1} \mathfrak{sl}(k_i))$ is the isotropy representation of $PL(l) \times H/PL(l) \times H_{<w>}$. The quotient space $(P/ker\rho, \theta)$ can be considered as a subbundle of P_tL, and we have the natural projection $\rho : P \to \widetilde{P}$. The group $PL(l)_{<v>}$ has the subgroup G_0 and the restriction of the isotropy representation ρ to $G_0 \times H_{<w>}$ is injective. The bundle P has the subbundle $h(P_tM) \times H$ with structure group $G_0 \times H_{<w>}$. Hence the restriction of $\rho : P \to \widetilde{P}$ to $h(P_tM) \times H$ is injective. Thus we obtain the sequence of reduction $(h(P_tM) \times H, \theta) \subset (\widetilde{P}, \theta) \subset P_tL$ corresponding to $G_0 \times H_{<w>} \subset \rho(PL(l) \times H_{<w>}) \subset GL(lk_1 \cdots k_{j-1} - k_j) \otimes GL(k_j)$. Hence L admits a reduction $h(P_tM) \times H$ of the Grassmannian structure P_tL of L.

7.2 Case of Lie groups

In the case of Lie groups the base spaces obtained by successive castling transformations are described more explicitly. Let ∇ be a projective structure on a n-dimensional Lie group L. Then we can construct a projective Cartan connection (Q, ω), where Q is a principal fiber bundle over M with structure group $PL(n+1)_v$, and we denote by h a injective bundle map from P_L to Q. Then by a successive castling transformations of (Q, ω) we obtain a Cartan connection $(Q \times \prod_{i=1}^{j} PL(k_i), \omega' \times \prod_{i=1}^{j} (\Lambda_1)_i)$ over a manifold N whose type is a subgeometry of the projective space $P(\boldsymbol{R}^{n+1} \otimes \otimes_{i=1}^{j} \boldsymbol{R}^{k_i})$. Then the base space is described as follows:

Proposition 7.2. *the base space N is isomorphic to the products $L \times \prod_{i=1}^{j} PL(k_i)$.*

Proof. The proof is similar to the one of Proposition 6.4. The base space N is diffeomorphic to the quotient manifold $Q \times \prod_{i=1}^{j} PL(k_i)/PL(l) \times \prod_{i=1}^{j} PL(k_i)_w$ and by Proposition 6.3 the isotropy group $PL(l) \times \prod_{i=1}^{j} PL(k_i)_w$ is given by the set $\{(f_0(g), f_1(g), \ldots, f_j(g)) \mid g \in PL(n+1)_v\}$, where $f_0 = id$ or $*$. Any element of Q is written by the form $h(x)g_0$ for some $x \in \mathcal{L}(L)$ and $g_0 \in PL(n+1)_v$. We denote by ρ a projection from Q to $\mathcal{L}(L)$. The frame bundle $\mathcal{L}(L)$ is isomorphic to the product $L \times GL(n)$ and an element $x \in \mathcal{L}(L)$ is written as $(a(x), A(x))$. Now we construct the map $N \to L \times \prod_{i=1}^{j} PL(k_i)$ defined by

$$[h(x)g_0, g_1, \ldots, g_j] \mapsto (a(x), g_1 f_1'(g_0)^{-1} f_1' \circ \imath(A(x))^{-1}, \ldots, g_j f_j'(g_0)^{-1} f_j' \circ \imath(A(x))^{-1}),$$

where $f_i' = f_i$ when $(\Lambda_1)_i = \Lambda_1$ and $f_i' = *f_i$ when $(\Lambda_1)_i = *\Lambda_1$. Then this map is well defined and a bundle isomorphism. □

Remark 7.3. The author could not determine the diffeo type of base spaces generally obtained by successive castling transformations.

The Cartan connection $(Q\times\prod_{i=1}^{j-1} PL(k_i), \omega'\times\prod_{i=1}^{j-1}(\Lambda_1)_i)$ can be extended to Grassmannian Cartan connection, and from the proof of 6.4 its base space K is also given by the product $L\times\prod_{i=1}^{j-1} PL(k_i)$. Thus the base space obtained by successive castling transformations admitting a Grassmannian structure is also given by a product Lie group. Furthermore if the given projective structure $[\nabla]$ on L is left invariant (resp. flat) under the group action of L, then a projective structure on N is also left invariant (resp. flat), and the Grassmannian Cartan connection over K is also left invariant (resp. flat).

8 A certain classification of manifolds obtained by successive castling transformations

Let j, l and α be positive natural numbers such that $\alpha \leq l - \alpha$. We consider the equation $(*) : (l-\alpha)\alpha + k_1^2 + k_2^2 + \cdots + k_j^2 - j = lk_1 \cdots k_j - 1$. We define a castling transformation for a set of positive natural numbers.

Definition 8.1. *Let l be a fixed integer such that $l \geq 3$. Let $(k_1, k_2, \ldots, k_j)$ be a set of integers. We define an integer by $k_i' := lk_1 \cdots k_{i-1}k_{i+1} \cdots k_j - k_i$ for $j \geq 2$, $1 \leq i \leq j$, and by $k_{j+1}' := lk_1 \cdot \cdots \cdot k_j - 1$. If $j = 1$, we define k_1' to be $l - k_1$. We call the set $(k_1, \ldots, k_{i-1}, k_i', k_{i+1}, \ldots, k_j)$ a castling transform of $(k_1, k_2, \ldots, k_j)$ at i-th position, and $(k_1, \ldots, k_j, k_{j+1}')$ a castling transform at $(j+1)$-th position. We call each k_i' and k_{j+1}' a number obtained by castling transform.*

If k_i is a positive natural number for $1 \leq i \leq j$ and $(k_1, \ldots, k_j)$ satisfies the equation $(*) : (l-\alpha)\alpha + k_1^2 + k_2^2 + \cdots + k_j^2 - j = lk_1 \cdots k_j - 1$, then castling transform at any position gives another solution $(k_1, \ldots, k_{i-1}, k_i', k_{i+1}, \ldots, k_j)$ for the equation $(*)$ and k_i' is a positive natural number again.

We observe that when $j = 1$, α gives a solution for the equation $(*)$: $(l-\alpha)\alpha + k_1^2 - lk_1 = 0$. We investigate the whole solutions of $(*)$ given by the successive castling transformations from α.

If we repeat a castling transformation for $(k_1, k_2, \ldots, k_j)$ at the same position twice, then we obtain the same set as $(k_1, k_2, \ldots, k_j)$. From now on we assume that successive castling transformations does not include this repetition. Namely if a set $\theta := (k_1, k_2, \ldots, k_j)$ is a castling transform of a set θ' at i-th position, then we only consider a castling transform of θ at m-th position with $m \neq i$. A sequence $\theta_1 \to \cdots \to \theta_n$ obtained by successive castling transformations from θ_1 is said to be reduced if the sequence does not contain any repetition of castling transformation. Moreover we identify $(k_1, k_2, \ldots, k_j, 1)$ with $(k_1, k_2, \ldots, k_j)$.

Lemma 8.2. *Let $(k_1, k_2, \ldots, k_j)$ be a set obtained by successive castling transformations from α. If k_j is a number obtained by the castling transform, then k_j is the unique largest number in $(k_1, k_2, \ldots, k_j)$.*

Proof. Let $\theta_1 \to \theta_2 \to \cdots \to \theta_n$ be a reduced sequence obtained by successive castling transformations from α, where $\theta_1 = \alpha$. The proof is by induction on $n \geq m \geq 2$. We express θ_m as $(h_1, h_2, \ldots, h_n)$. We assume that h_n is a number obtained by a castling transform of θ_{m-1} and h_n is the largest number in $\{h_i\}_{1 \leq i \leq n}$. Then θ_{m+1} is a castling transform of θ_m at i-th position with $i \neq n$. A new number κ of θ_{m+1} obtained by castling transform is written as (1) $lh_1 \cdots h_{i-1}h_{i+1} \cdots h_n - h_i$ or (2) $lh_1 \cdots h_n - 1$, and in both cases $\kappa > h_n$. In each case θ_{m+1} can be written as (1) $(h_1, \ldots, h_{i-1}, \kappa, h_{i+1}, \ldots, h_n)$ and (2) $(h_1, \ldots, h_n, \kappa)$ respectively. Hence

κ is the unique largest number in θ_{m+1}. When $m = 2$, θ_2 can be $(\alpha, l\alpha - 1)$ or $l - \alpha$. In both cases a number obtained by castling transform is the unique largest number in θ_2. Hence the induction proves the lemma. ☐

Proposition 8.3. *Let* $(k_1, k_2, \ldots, k_j)$ *be a set obtained by successive castling transformations from* α. *Then we have* $k_i \geq \alpha$ *for* $1 \leq i \leq j$.

Proof. Denote by $\theta_1 \to \cdots \to \theta_n$ a reduced sequence obtained by successive castling transformations from α, where $\theta_1 = \alpha$. Denote by κ_i $(i \geq 2)$ a new number of θ_i obtained by castling transform of θ_{i-1}. Set $\kappa_1 := \alpha$. For instance $\theta_2 = (\kappa_1, \kappa_2) = (\alpha, l\kappa_1 - 1)$, and $\theta_3 = (\kappa_1, \kappa_2, \kappa_3) = (\alpha, l\kappa_1 - 1, l\kappa_1\kappa_2 - 1)$ and $\theta_4 = (\kappa_1, \kappa_4, \kappa_3) = (\alpha, l\kappa_1\kappa_3 - \kappa_2, l\kappa_1\kappa_2 - 1)$. Thus we see that θ_m $(m \geq 2)$ can be written as $(\kappa_{i_1}, \cdots, \kappa_{i_n})$ where $n \leq m$ and some κ_{i_j} is equal to κ_m. Then κ_{m+1} $(m \geq 2)$ is equal to $l\kappa_{i_1} \cdots \kappa_{i_{s-1}}\kappa_{i_{s+1}} \cdots \kappa_{i_n} - \kappa_{i_s}$ for some s $(1 \leq s \leq n)$ with $i_s \neq m$ or $l\kappa_{i_1} \cdots \kappa_{i_n} - 1$ since successive castling transformations does not include a repetition. By Lemma 8.2 κ_m is the largest number in θ_m, thus $\kappa_{m+1} > \kappa_m$ for $m \geq 2$. By definition κ_2 can be $l - \alpha$ or $l\alpha - 1$. From assumption $l - \alpha \geq \alpha$. Thus $\kappa_2 \geq \alpha$. Hence $\alpha = \kappa_1 \leq \kappa_2 < \kappa_3 < \cdots < \kappa_n$, which proves the proposition. ☐

From the proof of this proposition, for any given reduced sequence $\theta_1 \to \cdots \to \theta_n$ of successive castling transformations, we have $\theta_i \neq \theta_j$ if $i \neq j$.

Proposition 8.4. *Let* j, l *and* α *be positive natural numbers such that* $j \geq 2$, $l \geq 3$ *and* $\alpha \leq l - \alpha$. *Let* k_i $(1 \leq i \leq j)$ *be natural numbers such that* $2 \leq k_1 \leq k_2 \leq \cdots \leq k_j$. Assume that we have the equality $(*) : (l - \alpha)\alpha + k_1^2 + k_2^2 + \cdots + k_j^2 - j = lk_1 \cdots k_j - 1$. Moreover assume that $k_j \geq \alpha$. Then we have $0 < lk_1k_2 \cdots k_{j-1} - k_j < k_j$.

Proof. Put $h_j := lk_1k_2 \cdots k_{j-1} - k_j$. Suppose that we have the equality $(*) : (l - \alpha)\alpha + k_1^2 + k_2^2 + \cdots + k_j^2 - j = lk_1 \cdots k_j - 1$ and the inequality $2 \leq k_1 \leq k_2 \leq \cdots \leq k_j$. Now we assume $lk_1k_2 \cdots k_{j-1} - k_j \leq 0$. Then $lk_1k_2 \cdots k_{j-1}k_j \leq k_j^2$. From $(*)$ we have $(l - \alpha)\alpha + 1 + \sum_{i=1}^{j} k_i^2 - j - k_j \leq 0$. Thus

$$\begin{aligned} 0 &\geq (l - \alpha)\alpha + 1 - j + k_1^2 + \cdots k_{j-1}^2 \\ &\geq (l - \alpha)\alpha + 1 - j + 4(j - 1) \\ &= 3j + (l - \alpha)\alpha - 3 \geq 3j - 1. \end{aligned}$$

Since $j \geq 2$, the last expression is greater than or equal to 0. This is a contradiction. Hence $0 < lk_1k_2 \cdots k_{j-1} - k_j$.

Next we divide the proof into the two cases: $\alpha = 1$ and $\alpha \geq 2$. Firstly we consider the case $\alpha \geq 2$. Then l must satisfy $l \geq 4$. Now we prove that if $\alpha \leq k_j \leq l2^{j-1} - \alpha$, and $2 \leq k_i \leq l2^{j-1} - \alpha$ for $1 \leq i \leq j - 1$, then we have $\sum_{i=1}^{j} {k_i}^2 - l\prod_{i=1}^{j} k_i \leq 4(j - 1) + \alpha^2 - l2^{j-1}\alpha$. We prove this by using the idea and technique of the proof of [SK, Lemma 2 in p.42]. Put $b := l2^{j-1} - \alpha$, and assume $\alpha \leq k_j \leq b$ and $2 \leq k_i \leq b$ for $1 \leq i \leq j - 1$. We put $f(k_1, \ldots, k_j) := \sum_{i=1}^{j} {k_i}^2 - l\prod_{i=1}^{j} k_i$. For $1 \leq \mu \leq j - 1$ we set

$$M_\mu^\alpha := f(\underbrace{2, \ldots, 2}_{\mu}, \underbrace{b, \cdots, b}_{j-\mu-1}, \alpha) \text{ and } M_\mu^b := f(\underbrace{2, \ldots, 2}_{\mu}, \underbrace{b, \cdots, b}_{j-\mu-1}, l2^{j-1} - \alpha).$$

Then we have $M_\mu^\alpha = 2^2\mu + (j-\mu-1)b^2 + \alpha^2 - l2^\mu b^{j-\mu-1}\alpha$, and $M_\mu^b = 2^2\mu + (j-\mu)b^2 - l2^\mu b^{j-\mu}$. Since $M_{j-1}^\alpha - M_\mu^\alpha = (j-\mu-1)(2^2-b^2) - l2^\mu\alpha(2^{j-\mu-1} - b^{j-\mu-1})$, we obtain

$$\begin{aligned} & \frac{M_{j-1}^\alpha - M_\mu^\alpha}{b-2} \geq -(j-\mu-1)(b+2) + l\alpha(j-\mu-1)2^{j-2} \\ = \ & (j-\mu-1)\{-b-2+l\alpha 2^{j-2}\} \geq 0. \end{aligned}$$

On the other hand since $M_\mu^\alpha - M_\mu^b = \alpha^2 - b^2 + l2^\mu b^{j-\mu-1}\{b-\alpha\}$, we obtain

$$\frac{M_\mu^\alpha - M_\mu^b}{b-\alpha} = -(\alpha+b) + l2^\mu b^{j-\mu-1}.$$

When $\mu = j-1$, this value is equal to zero. Thus we have $M_\mu^\alpha - M_\mu^b \geq 0$. Since f attains the maximum at the boundary points, we obtain the desired assertion $f \leq M_{j-1}^\alpha$.

Now from the equality $(*)$ we have

$$\begin{aligned} 0 &= (l-\alpha)\alpha - (j-1) + k_1^2 + k_2^2 + \cdots + k_j^2 - lk_1\cdots k_j \\ &\leq (l-\alpha)\alpha - (j-1) + 4(j-1) + \alpha^2 - l2^{j-1}\alpha \\ &= (l-\alpha)\alpha + 3(j-1) + \alpha(\alpha - l2^{j-1}) \\ &= (l-\alpha)\alpha + 3(j-1) + \alpha(\alpha - l) - \alpha l(2^{j-1}-1) \end{aligned}$$

Since $j \geq 2, l \geq 4, \alpha \geq 2$ the last expression is negative. However this is a contradiction. It follows that from the assumption $k_j \geq \alpha$ we obtain $k_j > l2^{j-1} - \alpha$.

From now on we consider the case $j = 2$. Firstly we show that $k_2 > k_1 + \frac{l}{2}$. Assume that $k_2 \leq k_1 + \frac{l}{2}$. Then we have

$$\begin{aligned} 0 &= (l-\alpha)\alpha - 1 + k_1^2 + k_2^2 - lk_1k_2 \\ &\leq (l-\alpha)\alpha - 1 + 2k_2^2 - l(k_2 - \frac{l}{2})k_2. \end{aligned}$$

Since $k_2 > 2l - \alpha$, we have

$$\begin{aligned} 0 &< (l-\alpha)\alpha - 1 + (2-l)(2l-\alpha)^2 + \frac{l^2}{2}(2l-\alpha) \\ &= (l-\alpha)\alpha - 1 + (2l-\alpha)\{-\frac{3}{2}l^2 - 2\alpha + l(4+\alpha)\}. \end{aligned}$$

About the part of this expression we have

$$-\frac{3}{2}l^2 - 2\alpha + l(4+\alpha) = l(-\frac{3}{2}l + 4) + \alpha(l-2).$$

Since $\alpha \leq \frac{l}{2}$, the last expression is less than or equal to $l(-l+3) \leq -l$. It follows that $(l-\alpha)\alpha + (2l-\alpha)(-l) < 0$. This is a contradiction. Therefore $k_2 > k_1 + \frac{l}{2}$.

Finally assume that $h_2 \geq k_2$. This condition is equivalent to $lk_1 \geq 2k_2$. Then from the equation $(*)$ we have

$$\begin{aligned}0 &= (l-\alpha)\alpha - 1 + k_1^2 + k_2^2 - lk_1k_2 \\ &\leq (l-\alpha)\alpha - 1 + k_1^2 + k_2^2 - 2k_2^2 \\ &= (l-\alpha)\alpha - 1 + k_1^2 - k_2^2 \\ &< (l-\alpha)\alpha - 1 + k_1^2 - (k_1 + \frac{l}{2})^2 \\ &\leq (l-\alpha)\alpha - 1 - (lk_1 + \frac{l^2}{4}) \\ &\leq \frac{l^2}{4} - \frac{l^2}{4} - 1 - lk_1 < 0.\end{aligned}$$

This is a contradiction. Hence $h_2 < k_2$, which is our assertion in this Proposition.

Next we consider the case $j \geq 3$. Now assume that $k_j \leq \frac{l}{4}(j-1)k_{j-1}$. Then we have

$$\begin{aligned}0 &= (l-\alpha)\alpha - (j-1) + k_1^2 + \cdots + k_j^2 - lk_1 \cdots k_j \\ &\leq (l-\alpha)\alpha - (j-1) + k_1^2 + \cdots + k_j^2 - \frac{4}{j-1}k_1 \cdots k_{j-2}k_j^2.\end{aligned}$$

Since we have $\frac{4}{j-1}k_1 \cdots k_{j-2} \geq \frac{4}{j-1}2^{j-2} \geq j+1$, this yields

$$(l-\alpha)\alpha - (j-1) + k_1^2 + \cdots + k_j^2 - \frac{4}{j-1}k_1 \cdots k_{j-2}k_j^2 \leq (l-\alpha)\alpha - (j-1) - k_j^2.$$

The inequality $k_j > l2^{j-1} - \alpha$ gives

$$\begin{aligned}0 &< (l-\alpha)\alpha - (j-1) - (l2^{j-1} - \alpha)^2 \\ &\leq \frac{l^2}{4} - (j-1) - (3l)^2 < 0.\end{aligned}$$

This is a contradiction. Hence we obtain $k_j > \frac{l}{4}(j-1)k_{j-1}$.

Finally suppose that $h_j \geq k_j$. This condition is equivalent to $k_j \leq \frac{l}{2}k_1 \cdots k_{j-1}$. Then combining this assumption with the equation $(*)$ yields

$$\begin{aligned}0 &= (l-\alpha)\alpha - (j-1) + k_1^2 + \cdots + k_j^2 - lk_1 \cdots k_j \\ &\leq (l-\alpha)\alpha - (j-1) + k_1^2 + \cdots + k_j^2 - 2k_j^2 \\ &< (l-\alpha)\alpha - (j-1) + k_1^2 + \cdots + k_{j-1}^2 - (\frac{l}{4})^2(j-1)^2k_{j-1}^2 \\ &\leq (l-\alpha)\alpha - (j-1) - (j-1)(\frac{l^2}{4^2}(j-1) - 1)k_{j-1}^2.\end{aligned}$$

Since $j \geq 3$ and $k_{j-1} \geq 2$, the last expression is less than or equal to

$$\begin{aligned}& (l-\alpha)\alpha - (j-1) - (l^2 - 8) \\ =\ & (l-\alpha)\alpha - (j-1) - (l^2 - 8) \\ \leq\ & (l-\alpha)\alpha - l^2 + 6 \\ =\ & -l(l-\alpha) - \alpha^2 + 6 < 0.\end{aligned}$$

This is a contradiction, which concludes $h_j < k_j$.

Now we consider the case $\alpha = 1$. In this case we have to also consider the case $l = 3$. We can prove the inequality $h_j < k_j$ by almost the same way as the case $\alpha \geq 2$. In the following we give the outline of the proof with emphasizing the difference between the cases $\alpha = 1$ and $\alpha \geq 2$.

Firstly if we assume $2 \leq k_i \leq l2^{j-1} - 2$ for $1 \leq i \leq j$, then from Lemma 2 of [SK, p. 42] we can directly obtain $\sum_{i=1}^{j} {k_i}^2 - l \prod_{i=1}^{j} {k_i}^2 \leq 4j - l2^j$. Combining this inequality with the equation $(*)$ implies a contradiction by a similar argument to the case $\alpha \geq 2$. Hence we obtain $k_j > l2^{j-1} - 2$. Now we divide the proof into the two cases $j = 2$ and $j \geq 3$. When $j = 2$, by using the inequality $k_j > 2l - 2$ we can prove $k_2 > k_1 + \frac{l}{3}$ similarly to the case $\alpha \geq 2$, but not $k_2 > k_1 + \frac{l}{2}$. Moreover by using $k_2 > k_1 + \frac{l}{3}$, we can obtain $h_2 < k_2$. When $j \geq 3$, by using the inequality $k_j > l2^{j-1} - 2$, we can obtain $k_j > \frac{l}{4}(j-1)k_{j-1}$. Moreover if we suppose $h_j \geq k_j$, then combining $k_j > \frac{l}{4}(j-1)k_{j-1}$ with the equation $(*)$ yields a contradiction. Thus we obtain $h_j < k_j$. □

The proof of Proposition 8.4 is a generalization of the one of [Kat, Lemma 7.3].

By Proposition 8.3 and 8.4 we obtain the following:

Theorem 8.5. *Let l and α be positive natural numbers such that $l \geq 3$ and $\alpha \leq l - \alpha$. Let $\theta = (k_1, k_2, \ldots, k_j)$ be the set of positive natural numbers. Then θ is obtained by a finite number of castling transformations from α if and only if θ gives the solutions of the equation $(*)$: $(l-\alpha)\alpha - (j-1) + k_1^2 + \cdots + k_j^2 - lk_1 \cdots k_j = 0$ and satisfies $k_i \geq \alpha$ for $1 \leq i \leq j$.*

In fact Proposition 8.4 implies a stronger result: if the positive natural numbers $\theta = (k_1, k_2, \ldots, k_j)$ gives a solution of the equation $(*)$ and moreover θ is not contained in the cube $C_{\alpha}^{j} = \{(x_1, \ldots, x_j) \mid |x_i| \leq \alpha - 1\}$, then θ is obtained by a finite number of castling transformations from the solution α of $(*)$ with $j = 1$.

Thus we obtain the following.

Proposition 8.6. *Assume that there exists no positive integer solution $(k_1, \ldots, k_j)$ of the equation $(*)$ satisfying $k_i \leq \alpha - 1$. Then any positive integer solution of $(*)$ is obtained by a finite number of castling transformations from the solution α.*

From many computations of the equation $(*)$, it seems that there exists no positive integer solution $\theta = (k_1, k_2, \ldots, k_j)$ such that θ is contained in the cube C_{α}^{j}. Hence we conjecture the following.

Conjecture. *There exists no positive integer solution $(k_1, \ldots, k_j)$ with $k_i \leq \alpha - 1$ for the equation $(*)$.*

This conjecture is true for $\alpha = 1, 2, 3$ or $j = 1$.

Now let (Q, ω) be a Grassmannian Cartan connection of type (β, α) over a manifold M, where we assume $l = \alpha + \beta \geq 3$ and $\alpha \leq \beta$. Denote by $S(l, \alpha)$ the set of positive natural number solutions of the equation $(*)$. We consider the tree T of Cartan connections obtained by successive castling transformations from (Q, ω). Each node is written as $(Q \times \prod_{i=1}^{j} PL(k_i), \omega' \times \prod_{i=1}^{j} (\Lambda_1)_i)$, where $\omega' = \omega$ or $*\omega$ and $(\Lambda_1)_i = \Lambda_1$ or $(\Lambda_1)_i = *\Lambda_1$. The model space of $(Q \times \prod_{i=}^{j} PL(k_i), \omega' \times \prod_{i=1}^{j} (\Lambda_1)_i)$ is $PL(l) \times \prod_{i=1}^{j} PL(k_i) / PL(l) \times \prod_{i=1}^{j} PL(k_i)_w$, where w is a point of $P(\boldsymbol{R}^l \otimes \otimes_{i=1}^{j} \boldsymbol{R}^{k_i})$. We define the map $\Phi : T \to S(l, \alpha)$ by $(Q \times \prod_{i=1}^{j} PL(k_i), \omega' \times$

$\prod_{i=1}^{j}(\Lambda_1)_i) \mapsto (k_1, \ldots, k_j)$. Each node of T induces a base space $M_{k_1\times\cdots\times k_j}$ by Proposition 6.4. Thus we obtain the map $\overline{\Phi}$ from the set of the base space of nodes in T to $S(l,\alpha)$ defined by $\overline{\Phi} : M_{k_1\times\cdots\times k_j} \mapsto (k_1, \ldots, k_j)$. Moreover $\overline{\Phi}$ induces the map Ψ from the set of the fiber $\prod_{i=1}^{j} PL(k_i)$ of base space of nodes in T to $S(l,\alpha)$. The map Ψ is bijective from Theorem 8.5. Thus we obtain the following.

Theorem 8.7. *There is a one-to-one correspondence between the set of the structure group* $\prod_{i=1}^{j} PL(k_i)$ *of the base spaces obtained by a finite number of castling transformations from* (Q,ω) *and the set of solutions* $(k_1, \ldots, k_j)$ *with* $k_i \geq \alpha$ *and* $j \geq 1$ *of the equation*

$$(*)\ \alpha\beta + k_1^2 + \cdots + k_j^2 - (j-1) - (\alpha+\beta)k_1\cdots k_j = 0.$$

Each solution $(k_1, \ldots, k_j)$ *corresponds to a manifold equipped with a projective structure, which is projectively flat if* (Q,ω) *is flat.*

From this theorem we obtain Theorem 1.1.

REMARK 8.8. A principal fiber bundle $L = N/PL(k_j)$ is equipped with a projective structure again if $lk_1\cdots k_{j-1} - k_j = 1$ and this manifold corresponds to a solution $(k_1, \cdots, k_{j-1})$ of the equation $(*)$: $\alpha\beta - (j-2) + k_1^2 + \cdots + k_{j-1}^2 - lk_1\cdots k_{j-1} = 0$. Indeed N corresponds to the solution $(k_1, \ldots, k_j)$ and $(k_1, \ldots, k_{j-1}, lk_1\cdots k_{j-1} - k_j)$ also gives a solution of $(*)$. If we have $lk_1\cdots k_{j-1} - k_j = 1$, then $(k_1, \cdots, k_{j-1}, 1)$ is a solution of $(*)$. Thus $(k_1, \cdots, k_{j-1})$ is a solution of $(*)$. If we have $lk_1\cdots k_{j-1} - k_j \neq 1$, then L admits a a Grassmannian structure of type $(lk_1\cdots k_{j-1} - k_j, k_j)$, whose corresponding Grassmannian Cartan connection is flat if (Q,ω) is flat.

REMARK 8.9. The equation $(*)$ $\alpha\beta + k_1^2 + \cdots + k_j^2 - (j-1) - (\alpha+\beta)k_1\cdots k_j = 0$ is a generalization of the equation $(**)$ $a^2 + k_1^2 + \cdots k_j^2 - j - 2ak_1\cdots k_j = 0$ which we obtained in our last paper [Kat, Theorem1.1]. Indeed put $\alpha := a-1$ and $\beta := a+1$ in $(*)$. Then we obtain $(**)$.

REMARK 8.10. The author expects that probably we can prove the map $\overline{\Phi}$ is bijective.

Acknowledgments

The author expresses his gratitude to Dr. Thomas Bruun Madsen for his encouragement. The author wishes to thank the Osaka City University Advanced Mathematical Institute and the King's College London for financial support and hospitality.

References

[Aga] Y. Agaoka. Invariant flat projective structures on homogeneous spaces. *Hokkaido Math. J.*, 11(2):125–172, 1982.

[GC] W. M. Goldman and D. Cooper. A 3-manifold with no real projective structure. *arXiv:* 1207.2007v1, 2012.

[Gol] W. M. Goldman. Locally homogeneous geometric manifolds. In *Proceedings of the International Congress of Mathematicians. Volume II*, pages 717–744, New Delhi, 2010. Hindustan Book Agency.

[Han] T. Hangan. Tensor-product tangent bundles. *Arch. Math.* (Basel), 19:436–440, 1968.

[Ish] T. Ishihara. On tensor-product structures and Grassmannian structures. *J. Math. Tokushima Univ.*, 4:1–17, 1970.

[Kat] H. Kato. Left invariant flat projective structures on lie groups and prehomogeneous vecctor spaces. *Hiroshima Math. J.*, 2012.

[KN] S. Kobayashi and T. Nagano. On projective connections. *J. Math. Mech.*, 13:215–235, 1964.

[MS] Y. Machida and H. Sato. Twistor theory of manifolds with Grassmannian structures. *Nagoya Math. J.*, 160:17–102, 2000.

[NS] K. Nomizu and T. Sasaki. *Affine differential geometry*, volume 111 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1994. Geometry of affine immersions.

[OT] V. Ovsienko and S. Tabachnikov. *Projective differential Geometry Old and New*. Cambridge university press, 2005.

[SK] M. Sato and T. Kimura. A classification of irreducible prehomogeneous vector spaces and their relative invariants. *Nagoya Math. J.*, 65:1–155, 1977.

[Tan1] N. Tanaka. Projective connections and projective transformations. *Nagoya Math. J.*, 12:1–24, 1957.

[Tan2] N. Tanaka. On the equivalence problems associated with a certain class of homogeneous spaces. *J. Math. Soc. Japan*, 17:103–139, 1965.

[Tan3] N. Tanaka. On non-degenerate real hypersurfaces, graded Lie algebras and Cartan connections. *Japan. J. Math.* (N.S.), 2(1):131–190, 1976.

Hironao Kato
Osaka City University Advanced Mathematical Institute, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585 Japan
Department of Mathematics, King's College London, Strand, London WC2R 2LS, United Kingdom.

E-mail: katohiro@sci.osaka-cu.ac.jp